

DATE FILED: 05/06/2009
DOCUMENT NO: 35

B8/B8 CIP

IN THE CANADIAN PATENT OFFICE

Examiner : M. Gillen
Applicant : Biogen, Inc.
Application No.: 374,378
Filed : April 1, 1981
For : DNA SEQUENCES, RECOMBINANT DNA
MOLECULES AND PROCESSES FOR PRODUCING HUMAN
FIBROBLAST INTERFERON-LIKE POLYPEPTIDES

AFFIDAVIT OF WALTER C. FIER

EXHIBITS 1-20

SUGANO EXHIBIT 1002
FIERS V. SUGANO
INTERFERENCE NO. 105,661

Walter Charles Cornelius **FIERS**

Date and place of birth: January 31, 1931, Leper, Belgium

Nationality: Belgian

Married: three children

CURRICULUM VITAE	
Opened	20 02
Désachetée le	
<i>[Signature]</i>	
Commissaire des brevets	
In presence of examiner	<i>[Signature]</i>
en présence de l'examinateur	<i>[Signature]</i>

1. Studies

High School: "Koninklijk Atheneum", Leper, 1949

University: Faculty of Agricultural Sciences, Ghent:

Engineer of Chemistry and Agricultural Sciences, 1954

"Agrégé" for Higher Education (Biochemistry), 1960

PhD, 1963

2. Scientific Career

1954 - 1955 Fellowship of the IWONL

1956 Assistant, Laboratory of Physiological Chemistry, Faculty of Medicine,
University of Ghent

Fellowship of the Danish Government (Danish-Belgian Cultural Agreements)

1957 - 1959 Research Assistant with the NFWO

1960 - 1961 Research Fellowship from the Rockefeller Foundation, New York, USA

1960 - 1961 Research Fellow in Biology, California Institute of Technology, Pasadena,
CA, USA

1960 - 1962 Senior Research Assistant with the NFWO

1962 Research Associate, Institute for Enzyme Research, University of Wisconsin,
Madison, WI, USA

1963 Assistant Professor at the Faculty of Agricultural Sciences, University of
Ghent, Belgium

1967 Associate Professor at the Faculty of Sciences, University of Ghent, Belgium
Responsible for the postgraduate course in Molecular Biology

1967 Director-Head of the Laboratory of Molecular Biology, University of Ghent,
Belgium

1969 Professor of Molecular Biology at the Faculty of Sciences, University of
Ghent, Belgium

3. Residences Abroad

Oct. 1, 1956 - Sept. 30, 1957

Carlsberg Laboratory, Copenhagen, Denmark (with Prof. Dr. H. Holter)

Oct. 1, 1960 - March 30, 1962

California Institute of Technology, Department of Biophysics, Pasadena, CA, USA
(with Prof. Dr. R.L. Sinsheimer)

April 1, 1962 - Oct. 1, 1962, and May 1, 1963 - June 7, 1963

Institute for Enzyme Research, University of Wisconsin, Madison, WI, USA (with
Prof. Dr. H.G. Khorana)

This is EXHIBIT FIERS-1
to
the Affidavit of Walter C. Fiers
sworn before me
this 4th day of November, 2001

Commissioner for Oath or Notary Public

4. Scientific Awards

- 1961 - Triannual Award "J.B. Van Helmon" (period 1958-1960) of the Royal Flemish Academy of Medicine of Belgium
- 1966 - Award of the Flemish Chemical Society (1964-1965)
- 1971 - Medal of the Société de Chimie Biologique de France
- 1975 - Award "Doctor A. De Leeuw - Damry - Bouriant" of the National Foundation for Scientific Research of Belgium for the period 1970-1975 (mathematical, physical and chemical sciences)
- 1976 - Francqui Award (Francqui Foundation, Belgium)
- 1978 - Doctor honoris causa, Catholic University of Leuven, Belgium
- 1980 - Jenkinson Memorial Lecture, University of Oxford, UK
- 1986 - Dr. Beijerinck Gold Medal for Virology, Royal Dutch Academy of Sciences, The Netherlands
- Rik & Nel Wouters Prize for Cancer Research, Belgium
- 1989 - Arnois - Baillet Latour Prize, Belgium
- Carlos J. Finlay Prize (UNESCO Prize for Microbiology, including Immunology, Molecular Biology and Genetics)
- 1990 - Personal title of "Baron" and hereditary nobility conferred by H.M. the King of Belgium
- 1991 - Robert Koch Prize (Robert Koch-Stiftung, Bonn, Germany)

5. Memberships, Offices and Committee Assignments

- 1966 - Elected to the Council of the Belgian Biochemical Society
- Elected as a member of the European Molecular Biology Organization (EMBO)
- 1969 - Member of the Advisory Board of the "European Journal of Biochemistry"
- 1970 - Visiting professor at the Catholic University of Leuven, Belgium
- 1971 - Member of the European Association for Cancer Research
- Member of the Editorial Board of "Natuur en Techniek", The Netherlands
- 1972 - Chairman of the research group "Oncoviruses" (formed on behalf of the Higher Council against Cancer), Department of Health, Belgium
- Member of the Editorial Board of "Intervirology"
- Guest member of the Dutch "Working Group on Nucleic Acids" (SON)
- 1973 - Corresponding member of the Royal Academy of Belgium, Class of Sciences
- 1974 - Organizer of the EMBO Workshop "Restriction Enzymes and DNA sequences", De Cirkel, Drongen, Belgium
- Member of the International Scientific Committee of the "International Institute of Cellular and Molecular Pathology" (ICP), Brussels, Belgium
- Member of the Scientific Board of the Department of Molecular Biology, ULB, Brussels, Belgium
- 1975 - Member of the Liaison Committee for Recombinant DNA Research of the European Science Foundation
- Member of the NFWO Commission for Biochemistry and Molecular Biology
- 1976 - Member of the Council for Medical Ethics, Foundation for Medical Scientific Research of Belgium
- 1977 - Member of the Editorial Board of "Gene"
- Member of the Scientific Council for Cancer Research of the ASLK
- Member of the Council of the European Molecular Biology Organization
- Member of the Overseas Advisory Panel of "The Biochemical Journal"

- 1978 - Second vice-president of the Belgian Biochemical Society
- 1979 - "Chaire Francqui" at the State University of Liège, Belgium
 - Member of the IUPAB Commission on Subcellular and Macromolecular Biophysics
 - Member of the Programme Committee for the 5th International Congress for Virology

- Member of the Editorial Board of "Biochimie"
 - Member of the Scientific Board of Biogen Inc.
- 1980 - Member of the Advisory Panel of the NATO Advanced Study Institutes
- 1981 - Member of the Royal Academy of Belgium, Class of Sciences
- 1982 - President of the Scientific Council for Cancer Research of the ASLK
 - Member of the Editorial Board of "The EMBO Journal"
 - President of the Belgian Biochemical Society
- 1983 - Member of the Editorial Board of "Nucleic Acids Research"
 - Member of the Editorial Board of "Anticancer Research"
 - Commander in the Order of Leopold
- 1984 - Member of the National Committee for Biochemistry
- 1985 - President of the NFWO Commission for Biochemistry and Molecular Biology (until 1990)
- 1987 - Member of the Commission for Biotechnology, Flemish Council for Science Policy
 - Member of the Editorial Board of "Biotherapy"
- 1988 - Member of the FEBS Fellowship Committee
 - Member of the Editorial Board of "The European Journal of Immunology"
 - Member of the Editorial Board of "Molecular Biology Reports"
 - Member of the Editorial Board of "Biotechnology Therapeutics"
 - Member of the Editorial Board of "Methods in Molecular and Cellular Biology"
 - Member of the Cell Board Subcommittee of the "Medical Research Council" (UK)
 - Member of the EEC Study Group on Ethical, Social and Legal Aspects of the Predictive Medicine Programme
- 1989 - Member of the Editorial Board of "Cytokine"
 - Member of the "Academia Europaea"
 - Elected member of "The Human Genome Organisation" (HUGO)
 - Member of the Board of the "Fondation van Gysel for Medical Research"
- 1990 - President of the Scientific Council for Cancer Research of the ASLK-Insurances
 - Corresponding member of the "American Association for Cancer Research"
 - Member of the Scientific Steering Committee for the EMBL (appointed by the Flemish Executive)
 - Belgian representative to the Council of the "International Society for Interferon Research" (ISIR)
 - Civil Cross First Class
- 1991 - Member of the "Scientific Advisory Committee" (SAC) for the EMBL (appointed by the EMBL Council)
 - Member of the Editorial Board of "Cancer Communications"
 - Honorary member of the Royal Flemish Society of Engineers
 - Member of the Scientific Board of the "Institut Pasteur du Brabant"
- 1992 - Member of the Editorial Board of "International Journal of Oncology"
 - Member of the Scientific Council of the "International Institute of Cellular and Molecular Pathology" (ICP), Brussels, Belgium
 - Honorary member of the Royal Academy of Medicine of Belgium
 - President of the "4th International Congress on Tumor Necrosis Factor and Related Cytokines", organized in Veldhoven, The Netherlands

"Chaire Francqui" at the Catholic University of Louvain (KUL), Faculty of Medicine, Belgium

1994 - *Member of the Editorial Board of "Circulatory Shock"*

1995 - *Member of the Editorial Board of "Lymphokine and Cytokine Research"*

- *Member of the Editorial Board of "Natural Immunity"*

- *Member of the Editorial Board of "The Journal of Inflammation"*

1996 - *Grand Officer in the Order of the Crown*

- *Retired as Professor at the University of Ghent and became Professor emeritus*

- *Director of the VIB (Flanders Interuniversity Institute for Biotechnology),
Department of Molecular Biology*

Dec 30 20 02
Walter FIERs

Commissioner of Patents
Commissaire des brevets

Office of the Examiner
Service de l'examinateur

LIST OF PUBLICATIONS

- A. Research papers
- B. Short communications
- C. a. Books (contributed chapters, editorships)
b. Reviews concerning own research
c. General (molecular biology, virology, etc.)
- D. Abstracts

This is EXHIBIT FIERs-2
to
the Affidavit of Walter C. Fiers
sworn before me
this 9th day of November, 2001

Commissioner for Oath or Notary Public
Board Assigned Page #121

A. RESEARCH PAPERS

- A.1 PIERCE, W. and STUCKE, J.
The chromatographic separation of mixtures of oligonucleotides and related products.
Biol. Soc. China, 45, 577-581, 1958.
- A.2 PIERCE, W. and WOLLEN, R. M.
A color method for the determination of ribonuclease activity.
Compt. Rend. Lab. Carlsberg 31, 561-566, 1960.
- A.3 PIERCE, W. and VANDERMEULEN, L.
Catalysis of nucleosides by barley extracts.
Arch. Intern. Physiol. Biochim. 18, 267-269, 1960.
- A.4 PIERCE, W.
The determination of ribonuclease activity.
Anal. Biochem. 3, 136-139, 1963.
- A.5 PIERCE, W. and VANDERMEULEN, L.
The ribonuclease-activity of barley.
Arch. Intern. Physiol. Biochim. 19, 339-343, 1963.
- A.6 PIERCE, W. and DE WACHTER, J.
The reaction mechanism of a nucleoside phosphorolysis.
Enzymologia 18, 199-210, 1963.
- A.7 PIERCE, W. and STEINBERGER, B. I.
The structure of the DNA of bacteriophage λ 174. I. The action of asopinucleosides.
J. Mol. Biol. 5, 498-519, 1963.
- A.8 PIERCE, W. and STEINBERGER, B. I.
The structure of the DNA of bacteriophage λ 174. II. Thermal inactivation.
J. Mol. Biol. 5, 420-433, 1963.
- A.9 PIERCE, W. and STEINBERGER, B. I.
The structure of the DNA of bacteriophage λ 174. III. Gistacitilogol evidence for a ring structure.
J. Mol. Biol. 5, 434-438, 1963.
- A.10 PIERCE, W.
An electron microscope for the imaging of ribonucleases, on topographical.
Verhandl. Ann. 91. Acad. Geneve. Belg. 25, 127-131, 1963.
- A.11 PIERCE, W. and STUCKE, J.
Studies on polynucleotides. XIII. Enzymic degradation. An exonuclease from *Lactobacillus acidophilus* 814. A. Purification, properties, and substrate specificity.
J. Biol. Chem. 238, 2380-2388, 1963.
- A.12 PIERCE, W. and BURRAN, W. B.
Studies on polynucleotides. XIII. Enzymic degradation. An exonuclease from *Lactobacillus acidophilus* 814. B. Stepwise degradation of oligonucleotides.
J. Biol. Chem. 238, 2389-2395, 1963.
- A.13 PIERCE, W., LIPKOWITZ, L. and VANDERMEULEN, L.
Studies on the bacteriophage λ 174. I. Distribution of purine sequences in the viral RNA and in yeast RNA.
J. Mol. Biol. 11, 411-419, 1963.
- A.14 PIERCE, W., DE WACHTER, J., LIPKOWITZ, L. and VANDERMEULEN, L.
Studies on bacteriophage λ 174. II. Mono-, di- and trinucleotide composition of the genomic ribonucleoside digest of λ 174 RNA.
J. Mol. Biol. 11, 431-441, 1963.
- A.15 DE WACHTER, J. and PIERCE, W.
Detection of uridine 3'-ethyl phosphate and nucleotides in digests of ribonucleic acid hydrolyzed with pancreatic ribonuclease in the presence of traces of ethanol.
Arch. Intern. Physiol. Biochim. 21, 141-144, 1963.
- A.16 PIERCE, W., VERLANGE, N. and VAN STEVENHAGEN, W.
The synthesis of bacteriophage λ 174 RNA in vitro. In Steinberg, R. V. and Busch, L. (eds.), Regulation of Nucleic Acid and Protein Biosynthesis, Elsevier Publishing Company, Amsterdam, pp. 118-166, 1963.
- A.17 DE WACHTER, J. and PIERCE, W.
External standardization in liquid scintillation counting of homogeneous acceptor labeled with one, two, or three isotopes.
Anal. Biochem. 10, 391-398, 1963.
- A.18 PIERCE, W.
Studies on the bacteriophage λ 174. III. Sedimentation heterogeneity of viral RNA preparations.
Virology 11, 413-424, 1963.
- A.19 DE WACHTER, J. and PIERCE, W.
Studies on the bacteriophage λ 174. IV. The 3'-end terminal undernucleotide sequence of the viral RNA chain.
J. Mol. Biol. 10, 307-313, 1963.
- A.20 WOLLEN, R., VAN MONTAGU, M. and PIERCE, W.
Model substrates for the rapid assay of specific ribonuclease activity.
Eur. J. Biochem. 2, 319-329, 1960.
- A.21 VANDERMEULEN, L., VAN STEVENHAGEN, W. and PIERCE, W.
Studies on the bacteriophage λ 174. V. The nucleoside 3'-triphosphate end groups of the copolymers of intracellular and the replicative form.
Eur. J. Biochem. 7, 114-123, 1966.

- A.22 MIN JOU, W. and PIERS, W.
Studies on the bacteriophage MS2. VII. Structure determination of the longer polynucleotide sequences present in the pancreatic ribonuclease digest of the viral RNA.
J. Mol. Biol. 40, 187-201, 1969.
- A.23 DE WACHTER, R. and PIERS, W.
Sequences at the 3'-terminus of bacteriophage Q β RNA.
Nature 221, 222-223, 1968.
- A.24 GILLIS, E., DE ROOVER, J.C. and PIERS, W.
The factor-dependence of bacteriophage Q β RNA-polymerase.
Arch. Intern. Physiol. Biochem. 72, 148-150, 1967.
- A.25 SLECHTA, M. and PIERS, W.
Studies on bacteriophage MS2. VIII. Evidence for direct formation of MS2 RNA by reaction with formaldehyde at acidic pH.
Biopolymers 2, 1333-1338, 1970.
- A.26 DE WACHTER, R. and PIERS, W.
Fractionation of RNA by electrophoresis on polyacrylamide gel slabs.
In Grossman, L. and Koidave, K. (eds.), "Nucleic Acids", Methods in Enzymology, Vol. 21, Part D, Academic Press, New York-London, pp. 147-178, 1971.
- A.27 CROSS, H.J., DUBINICK, F.B. and PIERS, W.C.
The RNA phosphopolymerase activity of bacteriophage MS2. A study on substrate specificity.
Eur. J. Biochem. 17, 118-123, 1970.
- A.28 NAEUMAN, G., MIN JOU, W. and PIERS, W.
Studies on the bacteriophage MS2. IX. The ribonucleotide sequences present in the pancreatic ribonuclease digest of the viral RNA.
J. Mol. Biol. 57, 597-613, 1971.
- A.29 DE WACHTER, R., VAN DERBEEK, A., NIEREGART, J., CONTREKAS, R. and PIERS, W.
The leader sequence from the 3'-terminus to the A-protein initiation codon in MS2-virus RNA.
Proc. Natl. Acad. Sci. USA 68, 281-289, 1971.
- A.30 DE WACHTER, R., NIEREGART, J., VAN DERBEEK, A., CONTREKAS, R. and PIERS, W.
Studies on the bacteriophage MS2. The untranslated 3'-terminal nucleotide sequence preceding the first start codon.
Eur. J. Biochem. 22, 409-414, 1971.
- A.31 DE WACHTER, R. and PIERS, W.
Preparative two-dimensional polyacrylamide gel electrophoresis of 32P-labeled RNA.
Anal. Biochem. 69, 181-187, 1972.
- A.32 MIN JOU, W., NAEUMAN, G., NIEREGART, R. and PIERS, W.
Nucleotide sequences of the gene coding for the bacteriophage MS2 coat protein.
Nature 237, 82-86, 1972.
- A.33 KIMMEL, E. and PIERS, W.
Studies on the bacteriophage MS2. X. The termination signal of the A protein cleavage.
J. Mol. Biol. 11, 217-241, 1972.
- A.34 VAN DERBEEK, A., NIEREGART, J., VAN DERBEEK, A. and PIERS, W.
Studies on the bacteriophage MS2. XI. Suppression of sensitive mutants of the A protein cleavage.
Mol. Gen. Genet. 119, 250-270, 1972.
- A.35 VAN DERBEEK, J. and PIERS, W.
Studies on the bacteriophage MS2. XII. Expansion of the virus in low salt.
Virology 50, 810-800, 1972.
- A.36 CONTREKAS, R., NIEREGART, J., MIN JOU, W. and PIERS, W.
Bacteriophage MS2 RNA: Nucleotide sequence of the end of the A protein gene and the intergenic region.
Nature New Biology 241, 99-101, 1973.
- A.37 SLECHTA, M. and PIERS, W.
Studies on the bacteriophage MS2. XIII. Conformation of MS2 RNA in acid medium.
Biopolymers 12, 1287-1291, 1973.
- A.38 SLECHTA, M. and PIERS, W.
Studies on the bacteriophage MS2. XIV. Fixation of the MS2 RNA acid structure by formaldehyde.
Biopolymers 12, 1293-1295, 1973.
- A.39 SLECHTA, M., CLAUWERT, J. and PIERS, W.
Studies on the bacteriophage MS2. XV. Hydrodynamic properties of the native and acid MS2 RNA structures.
Biopolymers 12, 1293-1295, 1973.
- A.40 NIEREGART, J. and PIERS, W.
Studies on the bacteriophage MS2. An internal nucleotide frequent recombining some ribosomal binding sites.
Eur. J. Biochem. 36, 119-121, 1973.
- A.41 VOLKHAUT, C. and PIERS, W.
A simple and highly sensitive method for sequence determination of 32P-labeled oligonucleotides.
Anal. Biochem. 62, 371-381, 1974.
- A.42 VAN DE WOODER, A., ROCHERS, A., VAN HEMERSCHE, J., VAN HEMERSCHE, M., VOLKHAUT, C. and PIERS, W.
Genomic nucleotide substitution: A new technique for sequence analysis of RNA.
Nucl. Acids Res. 1, 1025-1031, 1973.
- A.43 CONTREKAS, R. and PIERS, W.
A method for the isolation of cytoplasmic extracts from ribonuclease P1-oligonucleotides.
Anal. Biochem. 67, 119-128, 1972.
- A.44 VAN DERBEEK, A., VAN DE WOODER, A. and PIERS, W.
Location of the coat protein cleavage site in the MS2 RNA.
Virology 66, 240-243, 1973.

- A.43 VANDEBRIDGE, R., MIN JOU, W. and FIEBS, W.
3'-Terminal nucleotide sequence (n = 153) of bacteriophage MS2 RNA.
Proc. Natl. Acad. Sci. USA 73, 3528-3532, 1976.
- A.44 FIEBS, W., CONTRERAS, R., QUERINCK, F., NACZEMAN, G., VOLCHART, G., TIEBART, M., RAYMAKERS, A., WOLF, F. and VAN MONTAGU, M.
A-protein gene of bacteriophage MS2.
Nature 256, 213-216, 1975.
- A.45 DEVOIS, R., GILLIS, E. and FIEBS, W.
The carboxy addition of poly(A) to the 3'-end of RNA using bacteriophage MS2 RNA as a model system.
Eur. J. Biochem. 63, 401-410, 1976.
- A.46 YAMU, B.C., VAN DE VOORDE, A. and FIEBS, W.
Cleavage of the MS2-virus 49 genome by the restriction endonuclease III of *Neosporidia* sp.
Eur. J. Biochem. 63, 101-113, 1976.
- A.47 YANG, B.C., VAN DE VOORDE, A. and FIEBS, W.
Specific cleavage and physical mapping of MS2-virus RNA by the restriction endonuclease of *Neosporidia* sp.
Eur. J. Biochem. 63, 119-128, 1976.
- A.48 VOLCHART, G., MIN JOU, W. and FIEBS, W.
Analysis of 3'-labeled bacteriophage MS2 RNA by a mini-sequencing procedure.
Anal. Biochem. 71, 453-456, 1976.
- A.49 FIEBS, W., CONTRERAS, R., QUERINCK, F., NACZEMAN, G., VOLCHART, G., RAYMAKERS, A., VAN DEN BROECK, A., VOLCHART, G. and TIEBART, M.
Complete nucleotide sequence of bacteriophage MS2 RNA: Primary and secondary structure of the replicase gene.
Nature 268, 302-307, 1976.
- A.50 VAN ROY, F. and FIEBS, W.
Mycoplasma in African Green Monkey kidney cell cultures: Biochemical detection and effects in virus-infected cells.
In vitro 13, 737-763, 1977.
- A.51 DEVOIS, R., VAN EMMELO, J., QUERINCK-OPSTAMP, C., GILLIS, E. and FIEBS, W.
Addition of ATP RNA adenylyltransferase from *Saccharomyces cerevisiae* to 3'-linked oligo(A) to bacteriophage MS2 RNA and its effect on RNA replication.
Biochim. Biophys. Acta 447, 219-227, 1976.
- A.52 MIN JOU, W. and FIEBS, W.
Studies on the bacteriophage MS2. VIII. Comparison of the nucleotide sequence in related bacteriophage MS2s.
J. Mol. Biol. 103, 103-108, 1976.
- A.53 VAN DE VOORDE, A., CONTRERAS, R., ROCHERS, R. and FIEBS, W.
The initiation region of the MS2 RNA genome.
Cell 9, 117-120, 1976.
- A.54 VOLCHART, G., CONTRERAS, R., SOEDA, K., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of MS2 RNA virus 49 Hind III restriction fragment.
J. Mol. Biol. 110, 401-410, 1977.
- A.55 KERN, R. and FIEBS, W.
A two-dimensional electrophoretic procedure for the separation of DNA restriction fragments.
J. Mol. Biol. 110, 301-304, 1977.
- A.56 TIEBART, M., THYS, F., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the restriction fragments Hind III and Hind IV of MS2 RNA.
Nucl. Acids Res. 3, 1409-1422, 1976.
- A.57 MIN JOU, W., VAN MONTAGU, M. and FIEBS, W.
On the possible modulating role of the MS2 RNA-codon in bacteriophage MS2 RNA.
Biochem. Biophys. Res. Commun. 73, 1003-1007, 1976.
- A.58 ELTON, R.A. and FIEBS, W.
Rhythmic variations in purine run frequencies in bacteriophage MS2.
J. Theoret. Biol. 43, 49-58, 1976.
- A.59 DEVOIS, R., VAN EMMELO, J., GILLIS, E. and FIEBS, W.
Synthesis by avian-ovocellulose-virus RNA-dependent DNA polymerase of oligonucleotides complementary to bacteriophage MS2 polyomethylated in vitro.
Eur. J. Biochem. 73, 409-413, 1977.
- A.60 CONTRERAS, R., VOLCHART, G., THYS, F., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the restriction fragment Hind III of MS2 RNA.
Nucl. Acids Res. 3, 1003-1007, 1977.
- A.61 VAN MONTAGU, M., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the MS2 RNA restriction fragment Hind III - Map 2.
Nucl. Acids Res. 4, 1015-1018, 1977.
- A.62 VOLCHART, G. and FIEBS, W.
Micro thin-layer technique for rapid sequence analysis of 3'-labeled RNA: Double digestion and pancreatic ribonuclease analysis.
Anal. Biochem. 63, 178-181, 1977.

1985.
J. Mol. Biol. 104, 1047-1060, 1976.
- A.66 CONTRERAS, R., NOCIENS, R., VAN DE VOORDE, A. and FIEBS, W.
Overlapping of the vpi-vp3 gene and the vpi gene in the SV40 genome.
Cell 13, 339-346, 1977.
- A.67 MERRICAULT, J., VAN EMMELO, J., DEVOS, R., PORTER, A., FELLERS, P. and FIEBS, W.
The 3'-terminal nucleotide sequence of oncofeline sarcoma virus RNA.
Eur. J. Biochem. 83, 55-61, 1978.
- A.68 VERHEIJT, M., VAN MEUSEN, M., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of part of the Sialen virus 40 Hind-O restriction fragment. The presumed initiation region of the vpi gene.
Eur. J. Biochem. 83, 109-108, 1978.
- A.69 NOCIENS, R., VAN DE VOORDE, A., DEVOS, R. and FIEBS, W.
Nucleotide sequence of the Sialen virus 40 Hind-O restriction fragment.
Eur. J. Biochem. 83, 105-104, 1978.
- A.70 MARGENAU, G. and FIEBS, W.
Characterization of the 3'-terminal capped structures of late Sialen virus 40-specific mRNAs.
J. Virol. 33, 824-830, 1979.
- A.71 VAN ROY, P. and FIEBS, W.
Interference with Sialen virus 40 DNA replication by adenovirus type 3 during mixed infection of monkey cells.
J. Virol. 37, 373-381, 1979.
- A.72 CONTRERAS, R., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the restriction fragment Hind-O 3'-terminal of Sialen-virus-40 DNA (part of the vpi gene).
Eur. J. Biochem. 86, 317-324, 1978.
- A.73 VAN MEUSEN, M., VAN DE VOORDE, A. and FIEBS, W.
Complete nucleotide sequence of the Sialen-virus 40 Hind-O fragment and localization of the carboxyl terminus of the vpi protein.
Eur. J. Biochem. 86, 325-331, 1978.
- A.74 VAN MEUSEN, M., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the Sialen-virus-40 DNA region coding for the carboxyl-terminal part of the vpi antigen.
Eur. J. Biochem. 86, 333-344, 1978.
- A.75 PORTER, A.D., MERRICAULT, J., VAN EMMELO, J. and FIEBS, W.
Sequence of 135 nucleotides at the 3'-terminus of oncofeline sarcoma virus RNA.
Eur. J. Biochem. 87, 333-341, 1978.

- A.76 FIEBS, W., CONTRERAS, R., MARGENAU, G., NOCIENS, R., VAN DE VOORDE, A., VAN MEUSEN, M., MERRICAULT, J., VERHEIJT, M. and FIEBS, W.
Complete nucleotide sequence of SV40 DNA.
Nature 273, 117-120, 1978.
- A.77 VOLKERT, C., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the Sialen virus 40 Hind-O 3'-terminal region.
Proc. Natl. Acad. Sci. USA 75, 2180-2184, 1978.
- A.78 VAN ROY, P. and FIEBS, W.
Structure of the components of the adenovirus-Sialen virus 40 hybrid population adenovirus and Sialen virus 40 molecular interaction.
J. Mol. Biol. 116, 61-70, 1978.
- A.79 VAN MEUSEN, M., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the Sialen virus 40 Hind-O 3'-terminal region. The total amino acid sequence of the major structural protein vpi.
Eur. J. Biochem. 81, 413-420, 1978.
- A.80 VERHEIJT, M., VAN DE VOORDE, A. and FIEBS, W.
Nucleotide sequence of the Sialen virus 40 Hind-O 3'-terminal region. The total amino acid sequence of the late proteins VP3 and VP4.
Eur. J. Biochem. 81, 413-419, 1978.
- A.81 GROZINSKY, M., SANKOFF, D., MIN JOU, M., FIEBS, W. and FIEBS, W.
Characterization of the Sialen virus 40 Hind-O 3'-terminal region. A correlation between the stability of the coding antigen interaction and the choice of code words.
J. Mol. Biol. 117, 111-119, 1978.
- A.82 MARGENAU, G. and FIEBS, W.
Localization of the 3'-terminal of late SV40 mRNA.
Proc. Acad. Sci. USA 75, 2180-2184, 1978.
- A.83 DEVOS, R., VAN EMMELO, J., CONTRERAS, R. and FIEBS, W.
Construction and characterization of a plasmid containing a nearly full-size DNA copy of Sialen-virus-40 RNA.
J. Mol. Biol. 118, 575-619, 1979.
- A.84 DEVOS, R., CONTRERAS, R., VAN EMMELO, J. and FIEBS, W.
Identification of the translocatable element 131 in a Sialen-virus-40 DNA construct with plasmid pSV40 DNA in which a bacteriophage M13 DNA copy was inserted by the poly(dI)-poly(dI) linker method.
J. Mol. Biol. 118, 411-422, 1979.
- A.85 VOLKERT, C., FURTON, J., CRANFORD, L.V., DEAC, P. and FIEBS, W.
Nucleotide sequence of the coding region for small-t antigen of Sialen virus 40.
J. Virol. 30, 814-821, 1978.
- A.86 MIN JOU, M., MARGENAU, G. and FIEBS, W.
Characterization of Sialen-virus-40 RNA.
Eur. J. Biochem. 103, 589-595, 1978.

- A.87 HERRMANN, J., VAN DER VORDE, A. and PIERIS, W.
Nucleotide sequence of the Slon virus genome
J. Virol. 90, 182-186, 1979.
- A.88 VAN DER VORDE, A. and PIERIS, W.
Isolation and characterization of a specific deletion
mutant of tobacco etch virus type 3.
Virology 84, 484-490, 1979.
- A.89 MARGENAU, C., VAN NIEUWSTADT, D., CHRISTENSEN, D. and
PIERIS, W.
Stability of the 3' terminus of late mRNA induced
by a viable Slon virus to deletion mutant.
J. Virol. 31, 484-493, 1979.
- A.90 VAN NIEUWSTADT, D. and PIERIS, W.
Nucleotide sequence of the Hind-I fragment of Slon
virus to DNA. Comparison of the 5'-untranslated
region of wild-type virus and of above deletion
mutants.
Eur. J. Biochem. 100, 31-40, 1979.
- A.91 FORSLIN, K.E., ISERENTANT, D., CONTREARAS, R. and PIERIS,
W.
Polymorphic linker molecules for recombinant DNA
constructions.
Gene 7, 129-132, 1978.
- A.92 ISERENTANT, D. and PIERIS, W.
Secondary structure of the 3' end of bacteriophage
MS2 RNA: methylation and rational modification.
Eur. J. Biochem. 101, 595-604, 1979.
- A.93 WATTMAN, G., ISERENTANT, D., CHRISTENSEN, D. and PIERIS,
W.
Characterization of the major altered leader sequence
of late mRNA induced by SV40 deletion extent d1-101.
Nucl. Acids Res. 7, 1799-1814, 1979.
- A.94 ISERENTANT, D., VAN NIEUWSTADT, D. and PIERIS, W.
Studies on the bacteriophage MS2 RNA: status of the
early mutation.
J. Mol. Biol. 139, 243-261, 1980.
- A.95 ISERENTANT, D. and PIERIS, W.
Secondary structure of mRNA and efficiency of
translation initiation.
Gene 8, 1-12, 1980.
- A.96 VAN NIEUWSTADT, D. and PIERIS, W.
Recombination sequence for the restriction endonuclease
SfiI from Saccilar glabris.
Gene 8, 183-185, 1980.
- A.97 VOLKERTZ, G., VAN DE VOORDE, A. and PIERIS, W.
Nucleotide sequence of the Slon virus to Hindis +
III restriction fragment A second part of the 3'
antigen gene.
Eur. J. Biochem. 100, 149-157, 1980.
- A.98 VAN HERRMANN, J., VAN DE VOORDE, A. and PIERIS, W.
Nucleotide sequence of the Slon virus to Hindis +
III restriction fragment B fourth part of the 3'
antigen gene.
Eur. J. Biochem. 101, 157-167, 1980.
- A.99 VAN HERRMANN, J., VAN DE VOORDE, A. and PIERIS, W.
Nucleotide sequence of the Hindis+III restriction
fragment C third part of the 3' antigen gene) and
fourth part of the 3' antigen gene).
Eur. J. Biochem. 101, 167-176, 1980.
- A.100 VAN NIEUWSTADT, D., VAN DE VOORDE, A., VAN
HERRMANN, J., VOLKERTZ, G., DE WINTER, F. and
PIERIS, W.
Nucleotide sequence of Slon virus to Hindis + III restriction
fragment D third part of the 3' antigen gene) and
fourth part of the 3' antigen gene).
Eur. J. Biochem. 101, 177-186, 1980.
- A.101 MIN JOU, W., VERHOEVEN, E., DEVOS, E., SARAH, E.,
VOLKERTZ, G., PIERIS, W., THURILLALL, G.,
BARKER, C., CHAIK, W. and EXTERME, Y.
Complete structure of the noncoding region of the
Slon virus genome. The noncoding region contains a
polyoma-like sequence (PVS) (Herrmann and Min Jou)
Cell 19, 883-892, 1979.
- A.102 VAN DER VORDE, A., DEVOS, E., VERHOEVEN, E.,
VOLKERTZ, G., SARAH, E., PIERIS, W. and PIERIS, W.
Construction and characterization of a plasmid
containing a multiple cloning site derived from the
Yohann bacteriophage DNA copy of satellite
DNA.
J. Mol. Biol. 141, 159-171, 1980.
- A.103 VERHOEVEN, E., VAN DER VORDE, A. and PIERIS, W.
Total nucleotide sequence of a newly full-length
copy of satellite tobacco necrosis virus RNA.
J. Mol. Biol. 142, 117-127, 1980.
- A.104 HALPERIN, G. and PIERIS, W.
Characterization of the 3' terminal cop structure of
coxsackievirus B2.
J. Virol. 33, 255-261, 1980.
- A.105 VERHOEVEN, E., SARAH, E., MIN JOU, W., DEVOS, E.,
VOLKERTZ, G., SARAH, E. and PIERIS, W.
Antigenic drift between the homologous protein of the
Slon virus and the homologous protein of the
Victor virus.
Nature 286, 771-774, 1980.
- A.106 DENNY, R., COMPTON, J., DE CLERCQ, E., VOLKERTZ,
G., TAVELIER, J., DEVOS, E. and PIERIS, W.
Isolation and structure of a human simian
interferon gene.
Nature 281, 513-517, 1980.

- A.108 **CHEN, R., VAN DERBEEK, J. and FIER, W.**
Specific binding of interferon initiation factor 1 to
Satellite Tobacco Necrosis Virus RNA at a 1'-terminal
sequence comprising the ribosome binding site.
Proc. Natl. Acad. Sci. USA 78, 1311-1316, 1981.
- A.109 **CONTRERAS, R. and FIER, W.**
Initiation of transcription by RNA polymerase II in
coronavirus SV40-infected or noninfected, CV1 cells.
Mol. Acids Nuc. 9, 213-216, 1981.
- A.110 **THOMAS, J., DEWICK, R. and FIER, W.**
Evidence for a unique human fibroblast interferon
(IFN- β) chromosomal gene, devoid of intervening
sequences.
Mol. Acids Nuc. 9, 441-473, 1981.
- A.111 **RENAULT, E., STANISLAU, P. and FIER, W.**
Plasmid vectors for high-efficiency expression
controlled by the pL promoter of coliphage lambda.
Gene 15, 81-83, 1981.
- A.112 **VAN DER, P., FRANK, L. and FIER, W.**
Phosphorylation patterns of tumor antigens in cells
lytically infected or transfected by Sabin Virus 18.
J. Virol. 40, 24-34, 1981.
- A.113 **PANG, R., MIN JOU, W., RUTENBERG, D., DEVOS, R. and
FIER, W.**
Complete structure of 5' duck/duck/duck influenza
hemagglutinin gene: Animal virus as progenitor of
human H1 Hong Kong 1988 influenza hemagglutinin.
Cell 15, 313-323, 1981.
- A.114 **VAN ROPPE, L., MIN JOU, W., RUTENBERG, D., DEVOS, R.
and FIER, W.**
Complete nucleotide sequence of the nucleoprotein
gene from the human influenza strain A/WSN/33
(H3N2).
Eur. J. Biochem. 119, 347-353, 1981.
- A.115 **DEGRAVE, W., DEWICK, R., TAVERNIER, J., HUGENH, G.
and FIER, W.**
Nucleotide sequence of the chromosomal gene for human
fibroblast (H1) interferon and of the flanking
regions.
Gene 14, 137-143, 1981.
- A.116 **VOLCENT, C., TAVERNIER, J., DEWICK, R., DEVOS, R.
and FIER, W.**
Molecular mechanisms of nucleotide-sequence
rearrangements in cDNA clones of human fibroblast
interferon RNA.
Gene 15, 313-323, 1981.

- A.117 **CHEN, C., CHEN, D. and FIER, W.**
High-level synthesis in Escherichia coli of the SV40
small-t antigen under control of the bacteriophage
lambda pL promoter.
Gene 15, 43-54, 1981.
- A.118 **CHEN, C., ISHIGURO, D., DEWICK, R. and FIER, W.**
Systematic alteration of the nucleotide sequence
preceding the translation initiation codon and the
effects on bacterial expression of the cloned SV40
small-t antigen gene.
Gene 15, 55-63, 1981.
- A.119 **RENAULT, E., DE WILDE, P., MARMONNET, A., STANISLAU,
P. and FIER, W.**
Functional expression of individual plasmid-coded RNA
bacteriophage MS2 genes.
EMBO J. 1, 105-109, 1982.
- A.120 **KATZLEIN, R., RENAUD, E., FIER, W. and VAN DER,
J.**
Lysate gene expression of RNA phage MS2 depends on a
frameshift during translation of the overlapping coat
protein gene.
Nucleic Acids Res. 10, 33-41, 1982.
- A.121 **CHEN, C. and FIER, W.**
Expression and action of human fibroblast 41
interferon in monkey cells after transfection with a
recombinant SV40 plasmid vector.
J. Mol. Appl. Genet. 1, 183-194, 1982.
- A.122 **GROSJEAN, R. and FIER, W.**
Preferential codon usage in prokaryotic genes: The
optimal codon-anticodon interaction energy and the
selective codon usage in efficiently expressed genes.
Gene 18, 189-209, 1982.
- A.123 **CHEN, C., DE WILDE, P., RENAUD, E., DEWICK, R., DE
FIER, W. and FIER, W.**
Secretory proteins induced in human fibroblasts under
conditions used for the production of interferon β .
Proc. Natl. Acad. Sci. USA 79, 1788-1791, 1982.
- A.124 **DEVOS, R., CHEN, C., FIER, W., FIER, W. and FIER, W.**
Isolation and characterization of 17H-gamma RNA
derived from allogeneic induced human splenocytes.
J. Interferon Res. 2, 409-422, 1982.
- A.125 **DEVOS, R., CHEN, C., FIER, W., DEGRAVE, W., VAN
DERBEEK, J. and FIER, W.**
Molecular cloning of human lambda interferon cDNA and
its expression in eukaryotic cells.
Mol. Acids Nuc. 10, 1587-1591, 1982.
- A.126 **TAKA, Y., DEVOS, R., TAVERNIER, J., CHEN, C.,
CHEN, C. and FIER, W.**
Cloning and structure of the human lambda interferon
c chromosomal gene.
EMBO J. 1, 931-938, 1982.

- A.122 CONTRERAS, R., CHENYU, R., DEGRACE, W., and FIER, W.
Diaply, efficient in vitro synthesis of capped RNA
vector for direct expression of cloned eukaryotic
genes. *Nucl. Acids Res.* 10, 6153-6161, 1982.
- A.123 WOODE, A. and FIER, W.
Evidence for the direct involvement of DNA
replication origin in synthesis of late SV40 RNA.
Science 220, 308-310, 1982.
- A.124 VAN ROY, P., FRANK, L., and FIER, W.
Complete nucleotide sequence of a human influenza
neuroaminidase gene of subtype H2N1/Victoria/77/75.
J. Mol. Biol. 161, 3-11, 1982.
- A.125 VAN ROY, P., FRANK, L., and FIER, W.
Metabolic turnover of phosphoprotein sites in SV40
virus 20 large T antigen.
J. Virol. 81, 442-444, 1983.
- A.126 FRANK, L., VAN ROY, P., and FIER, W.
Change in gene expression and protein
phosphorylation in murine cells, transfected or
selectively infected with wild type and mutant SV40
viruses.
J. Biol. Chem. 258, 3316-3320, 1983.
- A.127 DE GROOT, G., DE WIT, L., VAN DE VORST, A., DE
WIT, M., and FIER, W.
Use of monoclonal antibodies to detect human
placental alkaline phosphatase.
Clin. Chem. 29, 113-119, 1983.
- A.128 DEHAUT, E., TIAO, X., and FIER, W.
Improved plasmid vector with a thermostable
expression and temperature-regulated runaway
replication.
Gene 22, 181-183, 1983.
- A.129 CHEN, D., VAN DE VORST, A., CONTRERAS, R., VAN
DER REEF, J., DE WIT, L., and FIER, W.
SV40 virus 20 antigen carrying attenuated deletions
in the 77-bp-pair repeat region.
J. Virol. 47, 1-14, 1983.
- A.130 SCHILL, J., DEWIT, L., VAN DER REEF, J., and
FIER, W.
Expression and characterization of the product of a
human tissue interferon gene in Chinese hamster
ovary cells.
Proc. Natl. Acad. Sci. USA 80, 6681-6686, 1983.
- A.131 POORE, J.E., GIBSON, M.A., COOPER, R.E., REISS,
C.E., BURROUGHS, E.P., FIER, W., and DE WIT, L.
Isolation of a human endothelial cell line which is
activated by virus and by human interferon.
J. Exp. Med. 157, 1119-1121, 1983.
- A.132 BARON, J., SUMSTON, T.C., BONGIA, P., and FIER, W.
Kodak, E. and FIER, W.
Codon usage and translation. In vivo nuclear level
measuring of the M2 coat protein sequence.
J. Biol. Chem. 258, 10007-10012, 1983.
- A.133 CHEN, D., DE WIT, L., TAVENIER, J., and FIER, W.
Human fibroblast interferon RNA transcripts of
different sizes in poly(A) positive induced cells.
Mol. Acids Res. 11, 1417-1418, 1983.
- A.134 VERHOEVEN, M., VAN ROY, P., and FIER, W.
Complete nucleotide sequence of the influenza B
virus NS1/77/75 virus haemagglutinin gene and
comparison with the H1N1/40 haemagglutinin.
Nucl. Acids Res. 11, 1507-1513, 1983.
- A.135 DEHAUT, E., SUMSTON, T., and FIER, W.
Inducible high level synthesis of mature human
fibroblast interferon in Escherichia coli.
Nucl. Acids Res. 11, 4677-4688, 1983.
- A.136 DEHAUT, E., TAVENIER, J., CHEN, D., and FIER, W.
Molecular cloning of human interferon 2 gene and its
expression in E. coli.
Nucl. Acids Res. 11, 1507-1513, 1983.
- A.137 DEGRACE, W., TAVENIER, J., RULINCK, P., and FIER, W.
Cloning and structure of the human interferon 2
chromosomal gene.
EMBO J. 2, 1149-1153, 1983.
- A.138 KILLY, V.K., FIER, W., and STOK, Y.D.
Cloned human interferon 2, but not interferon-3 or -
4, induces expression of HLA-DR determinants by total
suspended and specific leukemic cell lines.
J. Immunol. 131, 130-135, 1983.
- A.139 VAN ROY, P., FRANK, L., and FIER, W.
Protein kinase activity in immune complexes of
SV40 virus 20 large T antigen and transformation-
associated cellular protein.
Mol. Cell. Biol. 3, 277-279, 1983.
- A.140 FLEISCHMANN, M., FLEISCHMANN, C.M., and FIER, W.
Activation of interferon action by mixtures of
recombinant and endogenous human interferons.
Antivir. Res. 3, 311-316, 1983.
- A.141 CHEN, D., FIER, W., TAVENIER, J., and FIER, W.
Cloning of interferon 2 gene and its expression
in Chinese hamster ovary cells after simultaneous
transfection of human alpha and gamma
interferon.
J. Interferon Res. 3, 21-23, 1983.

- A.139 ERARD, P., CORTESE, P., SMITH, R.A., FIERS, W., CORIOLANO, R., and MARSHALL, M.
Characterization of cellular factors that induce the cytolytic activity and the expression of γ cell growth factor receptors of a T cell hybrid.
J. Exp. Med. 156, 388-397, 1982.
- A.140 SIMONS, C., REMAUT, E., ALLET, R., DEVOS, R., and FIERS, W.
High-level expression of human interferon genes in Escherichia coli under control of the pL promoter of bacteriophage lambda.
Gene 28, 33-44, 1981.
- A.141 HARMON, A., REMAUT, E., VAN BOOM, J., and FIERS, W.
Oligonucleotide directed mutagenesis: Selection of mutants by homologization of DNA-sequences.
Mol. Gen. Genet. 123, 126-133, 1982.
- A.142 VAN DER MEULEN, J., AMELIOT, P., PLATTICK, R., and FIERS, W.
Controlled synthesis of the coat protein of satellite tobacco necrosis virus in Escherichia coli.
Virology 120, 22-30, 1983.
- A.143 DEVOS, R., DEWES, C., SCHMIDT, R.J., VAN DER MEULEN, J., and FIERS, W.
Purification of recombinant glycosylated human gamma interferon expressed in transformed Chinese hamster ovary cells.
J. Interferon Res. 2, 381-389, 1981.
- A.144 COLLINS, T., KENNEDY, A.M., CLAYTON, C., FIERS, W., GIMBROW, M.A., BURKOFF, R.J., and PAPER, J.B.
Human erythroleukemia T lymphocyte interactions with vascular endothelium and fibroblasts: Role of adhesion and target cell molecular.
J. Immunol. 129, 1876-1885, 1982.
- A.145 COLLINS, T., KENNEDY, A.M., WALK, C.T., BERT, J.M., KAPLAN, D.J., FIERS, W., AULT, R.A., GIMBROW, M.A., STROMINGER, J.L., and PAPER, J.B.
Immune interferon activates multiple class-II major histocompatibility complex genes and the associated invariant chain gene in human endothelial cells and dermal fibroblasts.
Proc. Natl. Acad. Sci. USA 81, 4917-4921, 1984.
- A.146 ERU, J.D., CONTRERAS, R., GHEISSEN, D., KIM, J., and FIERS, W.
A system for dominant transformation and plasmid amplification in Saccharomyces cerevisiae.
Bio/Technology 3, 431-438, 1985.
- A.147 GROMBERG, A., HERNIM, R., and FIERS, W.
Interferon is a strong inducer of α -microglobulin but not of transferrin receptors of B6E1 cells.
Cell. Immunol. 93, 124-127, 1983.
- A.148 HARMON, A., MULLER, P., HARMON, A., TAVENIER, J., VAN DER MEULEN, J., KENNEDY, A.M., COLLIER, A., RUTSCHMANN, M., and FIERS, W.
Molecular cloning of mouse tumor necrosis factor cDNA and its subunit expression.
Mol. Acids Res. 12, 447-453, 1983.
- A.149 STANLEY, P., REMAUT, E., and FIERS, W.
Alterations upstream from the Shine-Dalgarno region and their effect on bacterial gene expression.
Gene 28, 215-222, 1983.
- A.150 HARMON, A., FRANK, L., TAVENIER, J., VAN DER MEULEN, J., TIZARD, R., KENNEDY, A.M., SCHMIDT, R.J., VAN VLIET, A., and FIERS, W.
Molecular cloning and expression of human tumor necrosis factor and comparison with mouse tumor necrosis factor.
Eur. J. Biochem. 132, 515-522, 1983.
- A.151 VAN DE VOORT, A., DE GROOT, C., DE WILDE, P., DE BRUIJN, M.E., POLITT, O., DE GROOT, J., VANDERKAM, D., and FIERS, W.
Screening of sera and their effect on cancer patients using a monoclonal antibody directed against human placental alkaline phosphatase.
Eur. J. Cancer Clin. Oncol. 11, 65-71, 1983.
- A.152 BACIONI, C., MCCABE, D., TAVENIER, J., and FIERS, W.
Binding of human tumor necrosis factor to high affinity receptors on B6E1 and lymphoblastoid cells sensitive to growth inhibition.
J. Biol. Chem. 258, 13395-13397, 1983.
- A.153 DEVOS, R., JAHNKE, B., VANDERKAM, D., and FIERS, W.
Recombinant interleukin 2 induces immunoglobulin secretion in Staphylococcus aureus Cowan strain 1 immunized cells.
Immunol. Lett. 11, 101-103, 1983.
- A.154 PARASKE, R., VERHEIJEN, J.L., ARONZANA-ESTRADA, F., ROTHMUT, S., MENCIN, M., RUIZ-MADIE, R., and FIERS, W.
Defective IFN- γ production in the human monocyte. Role of increased sensitivity to the suppressive effects of prostaglandin E.
J. Immunol. 133, 122-125, 1984.
- A.155 MITSURE, J.M., G.M., HERNIM, J.P., DEVOS, R., COLLIER, T., FIERS, W., and VANDERKAM, J.P.
Structural studies of the carbohydrate chains of human γ -interferon.
Eur. J. Biochem. 134, 491-497, 1983.

A.144 VAN DE WOUDE, A., TERREIN, R., DE BRUIJN, J., DE
WILG, S., VAN DER KROFT, D., and FIER, W.
The occurrence of human placental alkaline
phosphatase (ALP) in extracts of normal, benign and
malignant tissues of the female genital tract.
Tumour Biol. 6, 343-353, 1985.

A.145 ARIENKAMP-DEJONGE, Y., VERLUTTER, J.L. and FIER, W.
Interferon as macrophage-activating factor. III.
Proliferative effects of interferon- γ on the
interleukin 1 secretory potential of fresh or aged
human monocytes.
J. Immunol. 136, 1444-1448, 1985.

A.146 POREN, J.S., BIVIAQUA, M.P., MINORICE, D.L.,
LAPIERRE, L.A., FIER, W., and WISNIEWSKI, M.A.
Two distinct monokines, Interleukin 1 and tumor
necrosis factor, each independently induce
biogenesis and transient expression of the same
antigen on the surface of cultured human vascular
endothelial cells.
J. Immunol. 138, 1480-1487, 1986.

A.147 FRANKSON, L., VAN DER MEYDEN, J., RUTSCHMAYER, M. and
FIER, W.
Recombinant tumor necrosis factor: its effect and its
synergism with interferon- γ on a variety of normal
and transformed human cell lines.
Eur. J. Cancer Clin. Oncol. 22, 419-428, 1986.

A.148 STANISSE, P., PEXAUT, E. and FIER, W.
Inefficient transfection initiation causes premature
transcription termination in the lacZ gene.
Cell 44, 711-716, 1984.

A.149 COLLINS, P., LAPIERRE, L.A., FIER, W., STROMINGER, J.L. and POREN, J.S.
Recombinant human tumor necrosis factor increases
RNA levels and surface expression of HLA-A, B
antigens in vesicular endothelial cells and dermal
fibroblasts in vitro.
Proc. Natl. Acad. Sci. USA 83, 414-419, 1986.

A.150 ABUD-FIND, M., RANSEL, B.J., KELLEY, V.E., FIER, W. and STROM, J.R.
Natural killer cell responses to interferon.
Clin. Immunol. Immunopathol. 19, 144-154, 1984.

A.151 FRANKSON, L., RUTSCHMAYER, M.R., VAN DER MEYDEN, J. and FIER, W.
Recombinant tumor necrosis factor: Specific
specificity for a variety of human and murine
transformed cell lines.
Cell. Immunol. 100, 260-267, 1988.

A.152 STORPER, A.H., GUINAH, E., FIER, W. and FROBER, J.S.
Recombinant tumor necrosis factor and tumor
interferon act singly and in combination to
transform human vascular endothelial cell
monolayers.
Ann. N.Y. Acad. Sci. 457, 14-24, 1984.

A.153 C. RUGGERIO, V., YAMAMOTO, J., FIER, W. and BADLONT, J.
Induction of the synthesis of tumor necrosis factor
receptors by interferon- γ .
J. Immunol. 136, 1443-1450, 1986.

A.154 BIVIAQUA, M.P., POREN, J.S., MAJEAN, C.B., FIER, W., COYAK, M.A. and STROMINGER, M.A.
Recombinant tumor necrosis factor induces
plasmodium activity in cultured human vascular
endothelial cells: characterization and cooperation with the
action of interleukin 1.
Proc. Natl. Acad. Sci. USA 83, 6311-6317, 1986.

A.155 BALEWITZ, E.A., LEE, A., ALDAM, C., MCGOIR, E.,
THOMAS, J.A., TAVENNER, J., and FIER, W.
Human tumor necrosis factor treated with recombinant human
tumor necrosis factor alone or in combination with
interferon.
Cancer Res. 46, 1990-1995, 1986.

A.156 POREN, J.S., CHAMBERLAIN, M.A., LAPIERRE, L.A.,
TAVENNER, J., FIER, W., BOTHWELL, A. and SPRINGER, T.A.
Overlapping patterns of activation of human
endothelial cells by interleukin 1, tumor necrosis
factor, and immune interferon.
J. Immunol. 137, 1883-1888, 1986.

A.157 BROUHAERT, P.C., LEROUX-ROELS, E.E., CUYPER, V.,
TAVENNER, J., and FIER, W.
In vivo anti-tumor activity of recombinant human and
murine interferon alone and in combination with murine IFN- γ
genes, on a syngeneic murine melanoma.
Int. J. Cancer 36, 783-789, 1985.

A.158 MEETMAN, C., COMTE, P., VOLKERT, G., DERTNICK, R., TAVENNER, J., and FIER, W.
Structural analysis of the sequence coding for on
inducible 15-kDa protein in human fibroblasts.
Eur. J. Biochem. 155, 433-437, 1986.

A.159 VALCAREO, R., UNCE, Y., MONTIEL, J., STROMINGER, J.,
and FIER, W.
Purification, cryopreservation and preliminary X-ray
data of the bacteriophage MS2.
J. Mol. Biol. 120, 287-291, 1986.

A.160 VAN DER MEYDEN, J., FRANKSON, L. and FIER, W.
Treatment of murine interferon- γ -sensitive and
-resistant P815 leukemia cells with tumor necrosis
factor in combination with murine interferon- α /2 of
t.
J. Interferon Res. 8, 511-518, 1988.

- Board Assigned Page #131

- A.194 FOUFART, P., VANDERHAEGHE, P., CATHIER, S., VAN
SIEGHE, J., HAZENBACH, G., KRUS, V., FIERS, W. and
CHATEL, J.
A cell growth modulating and differentiating activity
of recombinant human 35-kD protein (HIF-2, HIF-2A,
HIF-2B).
Cancer Res. 47, 3319-3324, 1987.
- A.200 DEATHY, G.B., MAYLOR, M.B., FIERS, W. and BALMILL,
P.B.
DNA fragmentation and cytotoxicity caused by tumor
necrosis factor is enhanced by interferon- γ .
Eur. J. Immunol. 17, 689-693, 1987.
- A.201 GAO, J., VAN ROY, P., MESSIAEN, L., COSSETY, J.,
LIEBAUT, G., COOPMAN, P., FIERS, W. and NABET, M.
Pathology of tumours produced in syngeneic Fischer
rats by fibroblast-like cells before and after
transfection with oncogenes.
Path. Res. Pract. 183, 48-51, 1987.
- A.202 DEATHY, G.B., MAYLOR, M.B., FIERS, W. and BALMILL,
P.B.
The effect of recombinant human tumour necrosis
factor on growth and macromolecular synthesis of
human epithelial cells.
Exp. Cell Res. 170, 428-438, 1987.
- A.203 BALMILL, P.B., MARO, B.C., MOORE, T. and FIERS, W.
Therapeutic potential of tumor necrosis factor- α and
interferon in experimental human ovarian cancer.
Cancer Res. 47, 4753-4758, 1987.
- A.204 POGGIO, P., CARPANELLI, G., DI VITO, M., GIANNINI, M.,
POMILITI, G., FIERS, W., CRESPIER, I. and BELARZELLI,
F.
Nuclear magnetic resonance analysis of tumor necrosis
factor induced alterations of phospholipid
satellites and pH in Friend leukemia cell tumors and
fibrosarcoma in mice.
Cancer Res. 47, 4491-4499, 1987.
- A.205 CRESPIER, I., DELEA, F., TERNI, M., MARINO, S.,
ENGLES, S., BAUNY, C., FORTA, C., SORIA, J., FIERS,
W. and TAVENIER, J.
Tumor necrosis factor induced acute phase proteins in
rats.
J. Biol. Regul. Homeost. Agents 1, 171-176, 1987.
- A.206 FURT-JONES, S.A., FIERS, W. and POEHL, J.
Necrosis induced in 1 induction on human endothelial
cells and dermal fibroblasts.
J. Immunol. 139, 3317-3324, 1987.
- A.207 SALAMILL, P., GIBSON, M., BURKE, F., MAYLOR, M.,
TALBOT, D., PUGH, S., TAVENIER, J. and FIERS, W.
Evidence for tumor necrosis factor/cachectin
production in cancer.
Lancet ii, 8228-8232, 1987.
- A.208 LEWIS-REUTHER, A., FORTA, C., BAUNY, M., FORTA, S.,
VACCHER, P., TAVENIER, J., BAUNY, C., and FIERS,
W.
Structure of tumor necrosis factor by X-ray solution
scattering and gelation studies by single crystal
X-ray diffraction.
J. Mol. Biol. 187, 389-394, 1985.
- A.209 LAFITON, L.A., FIERS, W. and POEHL, J.
Three distinct classes of regulatory cytokines
control endothelial cell WBC antigen expression.
Interactions with tumor necrosis factor and interferon
the effects of tumor necrosis factor and lymphokine
free zones of endothelial cells and fibroblast
interferon.
J. Exp. Med. 167, 181-188, 1988.
- A.210 NABET, M., FIERS, W. and MORTU, B.J.
The induction of tumor necrosis factor
inhibits the tumorigenic action of TGF- β against on
established tumor in mice. In contrast,
immunologically dependent, and limited by severe
toxicity.
J. Exp. Med. 167, 1043-1048, 1988.
- A.211 MARO, B.C., DECONETTI, C., TAVENIER, J. and FIERS,
W.
Tumor necrosis factor effects of murine tumor
necrosis factor (TNF) and interferon- γ (IFN- γ)
in organ culture of B16 melanoma cells and
heart tissue.
Int. J. Cancer 47, 410-413, 1988.
- A.212 BARDET, R.L., ECKHART, A.M., DE BRUIJN, R.W., P.
FIERS, W. and JERRE, J.
Combination treatment of colon adenocarcinoma in rats
with tumor necrosis factor and the interferon inducer
2397.
J. Interferon Res. 8, 219-223, 1988.
- A.213 DE WAELE, P., FIERS, W., VAN DE VORDE, A., MICHARD,
P. and FIERS, W.
Expression in non-lymphoid cells of mouse recombinant
interferon- γ directed against the tumor marker
human placental alkaline phosphatase.
Eur. J. Biochem. 176, 217-221, 1988.
- A.214 TOFFI, M.C., COFFERT, J., CHIFFO, J., EDWARDS, J.,
BLANCHARD, S., GARNIER, J., FORTA, C., GIBSON, M.,
FIERS, W. and FIERS, W.
Cells for IFN- γ (IFN- γ) tumor necrosis factor, and
IL-1 are expressed at high levels in the organs of
normal individuals.
J. Immunol. 141, 1101-1104, 1988.
- A.215 SUPPES, P., STANLEY, R., VAN ROY, P. and FIERS, W.
Involvement of a protein protease in tumor necrosis
factor-mediated cytotoxicity.
Eur. J. Biochem. 178, 231-234, 1988.

- A-216 PRODETTI, E., DELARMONTE, F., COMPAGNON, D., DE VITO, M., MOORE, G., MOSA, J., SERRILLI, P., PIERS, W., and COHEN, I. Tumor necrosis factor (TNF) induces early morphologic and metabolic alterations in retinal endothelial cells and fibroblasts in mice.
Int. J. Cancer 43, 382-391, 1989.
- A-217 VAN HINBERGEN, W. B., KOSTER, T., VAN DEN BERG, E. A., PRINCE, M. M., PIERS, W., and DEWES, J. Tumor necrosis factor increases the production of plasminogen activator inhibitor in human endothelial cells in vitro and in vivo.
Blood 77, 1447-1453, 1991.
- A-218 ARIMURA-FRISCHOS, F., MOCHLIN, R. C., MILLER, F., PIERS, W., and VERLITZKE, J. L. Autocrine overproduction of tumor necrosis factor under the influence of interferon- γ upregulates HLA-DQ gene induction in human monocytes.
Proc. natl. Acad. Sci. USA 85, 4087-4091, 1988.
- A-219 MOSELMAN, R., NEUBURN, A., DUMONT, J. E., PIERS, W., and SALANO, P. Endocytic pathway of recombinant murine tumor necrosis factor in L-937 cells.
J. Immunol. 141, 3184-3188, 1989.
- A-220 MULLENBROEK, D., MARTENS, C., VERHOEVEN, M., LOPEL, C., BAETEMAN, A., NIU JOU, M., and PIERS, W. High-level transient expression of influenza virus proteins from a series of SV40 late and early replacement vectors.
Gene 88, 163-181, 1989.
- A-221 BEETA, J., SUKOWSKA, A., KOJ, A., HERING, T., RICHMOTO, Y., COMENT, J., PIERS, W., VAN DER MEIJT, J., and GAULDET, J. Different preparations of natural and recombinant human interleukin-6 (IL-6), IL-6-like activity stimulate acute phase protein synthesis and uptake of e-metabolizable acid by cultured rat hepatocytes.
Int. J. Biochem. 21, 341-344, 1989.
- A-222 COOPMAN, P., VAN ROY, F., DRACONETTI, C., CAO, J., PIERS, W., NUNZIATA, M., and BARRETT, M. Tumorigenicity, invasiveness and metastatic capability of rat-1 rat cells before and after transfection with bovine papilloma virus type 1 DNA.
Clin. Expt. Metastasis, 7, 49-61, 1989.
- A-223 GROTHEN, J., and PIERS, W. Regulation by the murine host of responsiveness toward various neoplastic cell lines, but not toward self, through adoptive transfer of a helper T lymphocyte clone with antitumor specificity.
Cancer Res. 49, 1671-1676, 1989.
- A-224 JARVIS, R., DEVOS, R., CURRIE, V., and PIERS, W. Purification of human interleukin-6 produced in Escherichia coli.
Gene 79, 183-196, 1989.
- A-225 ZOFFTS, P., BRANT, N., VAN ROY, F., and PIERS, W. In combination with interferon- γ , interleukin-6 normal, unfractionated serum and rat anti-CD3 monoclonal antibody.
Anticancer Res. 7, 147-152, 1987.
- A-226 TAVERNIER, J., DEVOS, R., VAN DER HEYDEN, J., HAUGHERT, G., RADDER, R., FAHNE, F., KAWAKITA, E., VAN HINBERGEN, J., COMERTS, R., and PIERS, W. Expression of human and murine interleukin-6 in embryonic systems.
Oncogene 8, 491-503, 1989.
- A-227 OMELAS, S. E., CAMPBELL, R. L., CARREL, D., CRIVELLO, C. J., COMERTS, R., PIERS, W., VAN DER HORST, C. A., M. J. J. Transformation of hepatocellular carcinomas with the oncogenic ability to reduce genes.
Cancer 70, 137-144, 1989.
- A-228 VANHAESBROECK, B., GROTHEN, J., and PIERS, W. An ongoing in vivo immune response affects the abundance and distribution of tyrosine-activated killer cell precursors, but does not influence their broad spectrum target specificity.
J. Immunol. 143, 2194-2201, 1989.
- A-229 LEHMANN, R., REMUT, E., and PIERS, W. Correlation between temperature-dependent cytoplasmic solubility and proteolytic cleavage of a heterodimeric protein in technical cells.
Gene 87, 97-106, 1988.
- A-230 DEVAULT, R., VANHAESBROECK, B., SUPPES, P., VAN ROY, F., and PIERS, W. Lithium chloride potentiates tumor necrosis factor-mediated cytotoxicity in vitro and in vivo.
Proc. Natl. Acad. Sci. USA 86, 2420-2423, 1989.
- A-231 MALIK, A. T., A., CRIVELLO, C. J., PIERS, W., and BARRETT, F. E. Pharmacological effects of tumour necrosis factor in experimental ovarian cancer.
Int. J. Cancer 41, 910-915, 1989.
- A-232 VAN ROY, F. N., COOPMAN, P., SUPPES, P., LISSAU, G., NUNZIATA, M., CAO, J., DRACONETTI, C. M., PIERS, W., and MALIK, M. Effect of oncoprotein transfection or passage in vivo on malignant phenotypes of rat cells.
Anticancer Res. 9, 133-144, 1989.
- A-233 VAN ROY, F., LISSAU, G., MARTEL, M., and PIERS, W. Partial transfection of human tumor cell lines showing defective interaction between unmutated p53 gene product and SV40 large-T antigen.
Oncogene 5, 207-210, 1989.

- A.214 TAVENIER, J., WARMOND, A., SAUDIN, R., HAUGIER, C., VAN OSTADE, F., and FIEBS, W.
Analysis of the structure-function relationship of tumor necrosis factor. Monoclonal chimeric for proteinase-inhibitor properties and epitope analysis. *J. Mol. Biol.* 111, 191-201, 1988.
- A.215 VANHAESBROECK, B., TIMMERHA, H.T. M., PROCK, C.J., VAN ROY, F., VAN DEN BERG, E.A., and FIEBS, W.
Modulation of cellular susceptibility to the cytotoxic/cytostatic action of tumor necrosis factor by adenovirus E1 gene expression in cell type. *Virology* 118, 167-168, 1988.
- A.216 VAN RINDEN, V.M. M., VAN DEN BERG, E.A., FIEBS, W., and DOOIJSTRA, C.
Tumor necrosis factor induces the production of proteinase-type plasminogen activator by human endothelial cells. *Blood* 75, 1981-1988, 1990.
- A.217 CHOMPFF, C., ADI, S., ZODER, M., MOISER, A., FIEBS, W., and WEISZ, R.
Search for mediators of the lipogenic effects of tumor necrosis factor: potential role for interleukin 6. *Cancer Res.* 50, 4333-4336, 1990.
- A.218 VAN ROY, F., MARTEL, M., VIERINCX, R., DEBART, R., FIEBS, W., and LECLETCQ, C.
Hormone sensitivity in vitro and in vivo of v-src-transfected MCF-7 cell derivatives. *Int. J. Cancer* 44, 522-524, 1989.
- A.219 CARRELL, D., JANSZENS, W., DEGRAVE, P., VAN DEN BROECK, C.A. M.J., KINCHOR, J.S., FIEBS, W., and CONTRERAS, R.
Heterologous gene expression by filamentous phage: secretion of human interleukin-6 by Aspergillus nidulans. *Gene* 94, 147-154, 1990.
- A.220 VANDEMAELE, P., BROOTEN, J., and FIEBS, W.
IL-6 acts synergistically on the IL-2 response of an autoreactive T-cell clone: synergism correlates with increased intracellular IL-2, but not with a modified IL-2 receptor expression. *Cell. Immunol.* 130, 18-21, 1990.
- A.221 VANDEMAELE, P., DECLEY, W., LIEBERT, C., and FIEBS, W.
Development of a simple, sensitive and specific bioassay for interleukin-6 based on the proliferation of PMH 1188 cells. Comparison with other bioassays for IL-6. *J. Immunol. Meth.* 135, 21-33, 1990.
- A.222 SUHYS, P., DEBART, R., DE VALCK, D., VANHAESBROECK, B., VAN ROY, F., and FIEBS, W.
Tumor necrosis factor mediated cytotoxicity is correlated with proteinase-inhibitor activity, but not with arachidonic acid release per se. *Eur. J. Biochem.* 193, 403-415, 1991.
- A.223 GUISSET, Y., TISON, B., VANHAESBROECK, J., VANHAESBROECK, B., NAEZENS, G., FIEBS, W., and CONTRERAS, R.
Production and purification of recombinant human interleukin-6 secreted by the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.* 198, 317-322, 1991.
- A.224 ALZAGA, M.-L., VANDEMAELE, P., DEBART, R., FLOQUIN, J., DESCHOT-LENGUEN, M., FLAMAND, V., MOISER, M., LIA, G., URBAIN, J., FIEBS, W., and GOLDMAN, M.
Cytotoxic enzyme syndrome induced by the IL-3-IL-11 anti-CD3 monoclonal antibody in mice: Prevention by high doses of methylprednisolone. *J. Immunol.* 146, 1166-1173, 1991.
- A.225 MAJOUVAN, V., NALLET, S.T. A., MEACER, A., FIEBS, W., and FIEBS, W.
Role of tumor necrosis factor in flavone acetic acid induced tumor vascular endothelial shutdown. *Cancer Res.* 50, 5337-5343, 1990.
- A.226 LIEBERT, C., VAN BROECK, B., BROUDERMAN, P., SHAM, A., and FIEBS, W.
Involvement of the liver, but not of IL-6, in IL-1-induced degradation to the lethal effects of tumor necrosis factor. *J. Immunol.* 145, 2635-2643, 1991.
- A.227 TAKAHASHI, H., BROUDERMAN, P., and FIEBS, W.
Induction of tolerance of tumor necrosis factor and antitumor activity of tumor necrosis factor in mice. *Cancer Res.* 51, 1344-1352, 1991.
- A.228 KUBOTA, R., MOSELMANN, M., FIEBS, W., DIMONT, J.E., and CALARO, P.
Tumor necrosis factor induces apoptosis (programmed cell death) in renal transplant cells in vitro. *Am. J. Pathol.* 135, 433-437, 1991.
- A.229 VANHAESBROECK, B., MARTEL, M., VAN ROY, F., CONTRERAS, R., and FIEBS, W.
Expression of the tumor necrosis factor gene in tumor cells correlates with reduced tumorigenicity and reduced invasiveness in vivo. *Cancer Res.* 51, 2219-2226, 1991.

- A.230 DEVAERT, M., DE POTTER, C., VANHAESBROECK, B., VAN ROY, P. and FLEIS, W.
Induction of intratumour cell infiltration and necrosis in mouse melanoma cells by the combined treatment with tumor necrosis factor and lithium chloride.
Am. J. Pathol. 138, 227-239, 1991.
- A.231 VAN BLAUDEL, S., LIBERT, C. and FLEIS, W.
Interleukin-6 enhances the expression of tumor necrosis factor receptors on hepatic cells and hepatocytoma.
Cytokine 3, 148-154, 1991.
- A.232 COMTESSA, B., CARRET, D., RINGHORN, J.R., VAN DEN HOFDEL, C.A., M.J.J. and FLEIS, W.
Efficient KHz-like processing of a glucanase-interleukin-6 fusion protein by hepatocellular carcinoma and secretion of mature interleukin-6.
Bio/Technology 9, 118-121, 1991.
- A.233 VAN OSTADE, K., TAVENIER, J., FRANGA, T. and FLEIS, W.
Localization of the active site of human tumor necrosis factor (TNF) by rotational analysis.
EMBO J. 10, 827-836, 1991.
- A.234 VANHAESBROECK, B., VAN BLAUDEL, S., LENAERTS, A., SUFFRS, P., DEVAERT, M., LUCAS, M., VAN ROY, P. and FLEIS, W.
Two discrete types of tumor necrosis factor-resistant cells derived from the same cell line.
Cancer Res. 51, 2469-2473, 1991.
- A.235 LIBERT, C., VAN BLAUDEL, S., BROUCKAERT, P. and FLEIS, W.
The influence of modulating substances on tumor necrosis factor and interleukin-6 levels after injection of ovine tumor necrosis factor or lipopolysaccharide in mice.
J. Immunother. 10, 327-335, 1991.
- A.236 VERMINKX, K., VANDEY, J., VANDEL, M., FLEIS, W. and VAN ROY, P.
Genetic manipulation of E-cadherin expression by epithelial tumor cells causes an invasion suppressor role.
Cell 66, 107-119, 1991.
- A.237 GRUMFIELD, C., BOWER, M., ROY, J., MOSER, A.H., FLEIS, W., DINARIELLO, C.A. and FLEIS, W.
Interleukin 2 inhibits stimulation of hepatic lipogenesis by tumor necrosis factor, interleukin 1, and interleukin 6 but not by interferon- α .
Cancer Res. 51, 3893-3897, 1991.
- A.238 DEVAERT, M., SCHULZE-DITTMER, K., VAN ROY, P. and FLEIS, W.
Lithium chloride potentiates tumor necrosis factor-induced and interleukin 1-induced cytokine and cytokine receptor expression.
Cytokine 3, 204-207, 1991.
- A.239 TAVENIER, J., DEVOS, K., COMTESSA, B., TAVENIER, J., VAN DER MEYDEN, J., FLEIS, W. and PLATTINCK, C.
A human high affinity interleukin-3 receptor (IL3H) is composed of an 165 specific α chain and a β chain shared with the receptor for GM-CSF.
Cell 64, 1173-1181, 1991.
- A.240 MARTEL, M.M., MEYERS, J., RINGHORN, J.R., DE SOUZA, C.R., VERMINKX, K., MOGILINS, A., FLEIS, W.C. and VAN ROY, P.M.
Down-regulation of E-cadherin expression in Madin Darby canine kidney (MDCK) cells inside tumor of nude mice.
Int. J. Cancer 47, 922-926, 1991.
- A.241 DEVOS, K., PLATTINCK, C., VAN DER MEYDEN, J., COMTESSA, B., VANHAESBROECK, J., FLEIS, W. and TAVENIER, J.
Molecular basis of a high affinity murine interleukin-3 receptor.
EMBO J. 10, 2122-2127, 1991.
- A.242 VAN DER MEYDEN, J., DEVOS, K., PLATTINCK, C., FACHE, J., FLEIS, W. and TAVENIER, J.
Characterization of the murine IL-3 receptor complex with the use of a panel of monoclonal antibodies: relationship to the murine IL-3 receptor.
J. Immunol. 147, 3413-3418, 1991.
- A.243 LANGIER, E.G., FLEIS, W. and VAN HAESBROECK, B.M.
Effects of tumor necrosis factor on prostacyclin production and the barrier function of human endothelial cell monolayers.
Arteriosclerosis Thrombosis 11, 872-881, 1991.
- A.244 FRANCO, L., JINCHIL, A., DEMOLIER, J., MOLENAAR, P., FLEIS, W. and COMTESSA, B.
The nucleotide sequence of a third cyclophilin-homologue gene from Saccharomyces cerevisiae.
Yeast 7, 917-923, 1991.
- A.245 VANHAESBROECK, B., DECLERCK, M., VERCAUTEREN, D., VAN DE COCK, M., GROOTEN, J., LUTSCHER, R., BROCKHAUS, M., LENAERTS, A. and FLEIS, W.
Functional characterization of the human tumor necrosis factor receptor p75 in a transfected rat mouse 3 cell hybridoma.
J. Exp. Med. 174, 1013-1024, 1991.
- A.246 DEMOLIER, J., FLEIS, W. and COMTESSA, B.
Efficient synthesis of secreted murine interleukin 3 by Saccharomyces cerevisiae: Influence of 1', untranscribed regions and codon usage.
Gene 151, 207-213, 1991.
- A.247 FLEIS, W., DINARIELLO, C.A., FLEIS, W. and GRUMFIELD, C.
Stimulation of lipolysis in cultured fat cells by tumor necrosis factor, interleukin-1, and the interferon is blocked by inhibition of prostaglandin synthesis.
Endocrinology 110, 10-14, 1991.

- Board Assigned Page #136

- A.278 TAKAHASHI, K., BROUCKHAERT, P. and FIERS, W.
Cytotoxic agents inhibit the induction of
tolerance to the toxic effects of tumor necrosis
factor.
J. Immunother. 14, 18-21, 1991.
- A.280 DEWOLFE, J., DE BACHER, M., FIERS, W. and CORREIA, A.
Phenotypic effects in Saccharomyces cerevisiae after
regulated expression of 8-(1,2)-glucanase from
Bacillus pumilus.
J. Biotechnol. 21, 283-285, 1991.
- A.281 VANHAEREN, D., REED, J.C., DE VALCKE, D., GROOTEN, J.,
WITTEKAMP, R., TAKAZA, S., BEYAERT, R., VAN ROY, F.
Effect of bcl-2 proto-oncogene expression on cellular
sensitivity to tumor necrosis factor-mediated
cytotoxicity.
Oncogene 8, 1073-1081, 1992.
- A.282 STEIGLER, L., SEMAAT, E. and FIERS, W.
Lead as a carrier molecule for the functional
exposition of IgG-binding domains of the Staphylococcus
aureus protein A at the surface of Escherichia coli
cells.
Mol. Gen. Genet. 256, 187-192, 1991.
- A.283 DE VALCKE, D., BEYAERT, R., VAN ROY, F. and FIERS, W.
Tumor necrosis factor cytotoxicity is associated with
phospholipase D activation.
Eur. J. Biochem. 211, 491-493, 1991.
- A.284 VAN OSTADE, S., VANHAEREN, D., EVERAERT, S.,
LOOTSCHER, M., GENT, S., BROCKMUS, M., LESLAUER, M.,
TAVENIER, J., BROUCKHAERT, P. and FIERS, W.
Human TGF mutants with selective activity on the p33
receptor.
Nature 351, 368-369, 1992.
- A.285 GERARD, C., BRUNS, C., NACHMAY, A., ARAMONIC, D.,
VANHAEREN, D., DELVAUX, A., FIERS, W., GOLDMAN, M.
and FELD, Y.
Interleukin 1 α reduces the release of tumor necrosis
factor and prevents lethality in experimental
endotoxemia.
J. Exp. Med. 177, 547-550, 1992.
- A.286 DEVOS, S., GUISSE, E., CORREIA, S., VERHEE, A., VAN
DER MEYDE, J., KAMBERG, M., LAMM, M.W., FIERS, W.,
TAVENIER, J. and PLATTING, G.
Recombinant soluble human interleukin-3 (IL-3)
receptor molecules. Cross-linking and steric hindrance of
binding to IL-3.
J. Biol. Chem. 267, 6361-6367, 1992.
- A.287 GUISSE, E., HUBBARD, J., SEMAAT, E. and FIERS, W.
Binding of the p33 coat protein in Escherichia coli is
modulated by translational pauses resulting from mRNA
secondary structure and codon usage: A hypothesis.
J. Theor. Biol. 162, 261-272, 1991.
- A.288 BEYAERT, R., VANHAEREN, D., MEYER, P., MEYER, P., ZOOMER, E.,
DE VALCKE, D., SCHULZ-OSTHOFF, K., MACCEAN, G.,
VAN ROY, F. and FIERS, W.
Constitution of tumor necrosis factor cytotoxicity by
action of the protein kinase inhibitor staurosporine.
Cancer Res. 51, 2421-2430, 1991.
- A.289 BEYAERT, R., MEYER, P., DE VALCKE, D., MEYER, P.,
VAN ROY, F. and FIERS, W.
Enhancement of tumor necrosis factor cytotoxicity by
lithium chloride is associated with increased inositol
phosphate accumulation.
J. Immunol. 151, 391-396, 1993.
- A.290 VANHAEREN, D., ARAMONIC, D., MEYER, P., VAN DER
MEYDE, J., GROOTEN, J., GROOTEN, P., MOONEN-PELZ, E.L.,
GOLDMAN, M. and FIERS, W.
Increased IL-1 production and IL-1-mediated B α
secretion in murine bone-marrow-derived dendritic cells.
J. Immunol. 150, 4179-4183, 1993.
- A.291 GUISSE, E., DEMISEL, J., MEYER, P., VANHAEREN, D.,
PLATTING, G., HUBBARD, J., GROOTEN, P., MOONEN-PELZ, E.L.,
GOLDMAN, M. and FIERS, W.
High-level expression, purification and renaturation of
recombinant murine interleukin-3 from Escherichia coli.
Protein Expression Purification 4, 110-116, 1991.
- A.292 SCHULZ-OSTHOFF, K., BEYAERT, R., VANHAEREN, D.,
MACCEAN, G. and FIERS, W.
Regulation of the mitochondrial electron transport
chain by the cytotoxic and gene-inductive effects of
TNF.
EMBO J. 12, 1095-1104, 1993.
- A.293 GROOTEN, J., CORREIA, S., VANHAEREN, D. and
FIERS, W.
Cell membrane permeabilization and cellular collapse,
followed by loss of dephosphorylation activity: Early
events in tumor necrosis factor-induced cytotoxicity.
Cytokine 5, 318-325, 1993.
- A.294 STEDLER, L., SEMAAT, E. and FIERS, W.
p33 p11 as a vector system for surface exposition of
an immunoglobulin G-binding domain of protein A as
staphylococcus aureus in Escherichia coli.
J. Bacteriol. 175, 1639-1643, 1993.
- A.295 PATRICK, M.P., MACCEAN, G., VANHAEREN, D. and
FIERS, W.
Activation of the nuclear factor κ B is not sufficient
for regulation of tumor necrosis factor-induced
interleukin-4 gene expression.
Biochimica 13, 1007-1014, 1993.
- A.296 SLOAN, M.P., DE VALCKE, D., FIERS, W. and POKER, J.S.
Tumor necrosis factor activates human endothelial cells
through the p33 tumor necrosis factor receptor but the
p33 receptor contributes to activation of low tumor
necrosis factor concentration.
Am. J. Pathol. 141, 1136-1140, 1993.

- [illegible]

Abstracts of papers presented at the 1993 Annual Meeting of the Society for Experimental Biology and Medicine, 1993, 164, 1-151, 1993.

- A.313 DEHAYE, S., VANHAESBROECK, B., FLEIS, M., WILLIAMS, J., and JONIAU, M.
Protection by all-acid glycosylated against tumor necrosis factor-induced lethality.
J. Leukocyte Biol. 51, 141-151, 1993.
- A.314 MERTENS, M., REMAUT, M., and FLEIS, M.
Tight transcriptional control mechanism ensures stable high-level expression from T1 promoter-based expression plasmids.
BioTechnology 11, 175-179, 1993.
- A.315 LOS, M., VAN DE CRAEN, M., PENDING, L.C., SCHENK, H., WESTENDORP, M., BARNHART, P.A., BROCK, W., KRAMER, P.H., FLEIS, M., and SCHULIS-OSTHOFF, K.
Requirement of an ICS/CEB-3 protease for Fas/APO-1-mediated apoptosis.
Nature 365, 81-83, 1993.
- A.316 TAKAHASHI, M., BROUHAERT, P., and FLEIS, M.
Mechanism of tolerance to tumor necrosis factor: Receptor-specific pathway and selectivity.
Am. J. Physiol. 269, R188-R193, 1993.
- A.317 KOOZEK, V., BROUHAERT, P., DE VOS, K., and FLEIS, M.
Direct evidence for tumor necrosis factor-induced mitochondrial reactive oxygen intermediates and their involvement in cytotoxicity.
Proc. Natl. Acad. Sci. USA 92, 5115-5118, 1993.
- A.318 VANLANUSCHOT, P., BRIEMANT, S., DE VOS, K., and FLEIS, M.
Receptor-specific pathway and selectivity.
Am. J. Physiol. 269, R188-R193, 1993.
- A.319 VERLUMMEN, B., VANHAESBROECK, P., DECLERQ, W., VAN DE CRAEN, M., BROUHAERT, P., and FLEIS, M.
A fairly conserved epitope on the homocysteine of influenza A (H1N1) virus with variable accessibility to neutralizing antibody.
Virology 191, 516-518, 1993.
- A.320 VERLUMMEN, B., VANHAESBROECK, P., DECLERQ, W., VAN DE CRAEN, M., BROUHAERT, P., and FLEIS, M.
Influence of receptor-specific pathway and selectivity.
Am. J. Physiol. 269, R188-R193, 1993.
- A.321 VERLUMMEN, B., VANHAESBROECK, P., DECLERQ, W., VAN DE CRAEN, M., BROUHAERT, P., and FLEIS, M.
Influence of receptor-specific pathway and selectivity.
Am. J. Physiol. 269, R188-R193, 1993.
- A.322 FLEIS, M., REMAUT, M., and FLEIS, M.
Production in recombinant cell of a functional active and murine: human chimeric Fas-12 fragments and active antibody directed against human placental alkaline phosphatase.
J. Biotechnology 12, 111-113, 1993.
- A.323 MERTENS, M., REMAUT, M., and FLEIS, M.
Production of soluble and active recombinant murine interleukin-1 in recombinant cells: High level expression, titration-induced release, and purification.
Protein Expression Purification 5, 461-466, 1993.
- A.324 DECLERQ, W., VANHAESBROECK, B., VANHAESBROECK, P., BROUHAERT, P., and FLEIS, M.
Generation and biological characterization of monoclonal, unclonable murine tumor necrosis factor.
J. Biol. Chem. 268, 18471-18476, 1993.
- A.325 CORNELIS, S., FACHE, L., VAN DER HEYDEN, J., GUILLI, Y., TAVENIER, J., DEVOS, M., FLEIS, M., and PLATTICK, C.
Phosphorylation of critical residues in the cytoplasmic domain of the human interleukin-3 receptor α chain required for growth against transduction.
Eur. J. Immunol. 23, 1853-1866, 1993.
- A.326 BEAULT, R., VANHAESBROECK, B., DECLERQ, W., VAN DER HEYDEN, J., TAVENIER, J., DEVOS, M., FLEIS, M., and PLATTICK, C.
Cytosolic kinase phosphorylates the p135 tumor necrosis factor receptor and negatively regulates tumor necrosis factor signaling for apoptosis.
J. Biol. Chem. 268, 11193-11198, 1993.
- A.327 TAKAHASHI, M., FLEIS, M., and BROUHAERT, P.
Anti-tumor activity of tumor necrosis factor in combination with interferon- γ is not affected by prior tolerization.
Int. J. Cancer 53, 866-870, 1993.
- A.328 DECLERQ, W., VANHAESBROECK, P., and FLEIS, M.
Detection of chimeric Fas-12 fragments in the tumor necrosis factor (TNF) receptor transduces TNF signal: Necessity for the 135 kDa TNF receptor transmembrane domain.
Cytokine 7, 701-709, 1993.
- A.329 MERTENS, M., REMAUT, M., and FLEIS, M.
Vascular, multi-targeted plasmids for high-level expression of heterologous genes in recombinant cells: Overproduction of human and murine cytokines.
Gene 164, 9-15, 1993.
- A.330 STEINER, L., FLEIS, M., and REMAUT, M.
Stimulus-induced light activation of a chimeric Fas-12 fragment and expression of Fas-12 protein in recombinant cells: Dependence on using a single chimeric epitope.
BioTechnology 11, 1193-1195, 1993.

- A.311 COOSEN, V., GROUPE, J. and PIER, W.
The oxidative metabolism of glutamine. A modulator of reactive oxygen intermediates-mediated cytotoxicity of tumor necrosis factor in L1210 fibrosarcoma cells.
J. Biol. Chem. 271, 121-126, 1996.
- A.312 BERTHET, R., CUNHA, A., VAN DER MEYDEN, J., FLAISANCE, S., LEE, J.-C., HEDGECOCK, C., CONN, P. and PIER, W.
The p15^{INK4A} mitogen-activated protein kinase pathway regulates interferon- γ synthesis in response to tumor necrosis factor.
EMBO J. 15, 1918-1923, 1996.
- A.313 ELZOO, T., MIN JOU, W. and PIER, W.
Recombinant neuraminidase vaccine protects against lethal influenza.
Vaccine 14, 541-549, 1996.
- A.314 CAWLEY, A., PIER, W. and BROUHAERT, P.
Murine IL-12 is involved in endotoxin-Gal α 1n bacillus-induced sensitization and is by itself sufficient to sensitize mice to the lethal effects of human TNF.
J. Immunol. 156, 4688-4698, 1996.
- A.315 LESTER, C., VAN NELLE, W., BROUHAERT, P. and PIER, W.
Platelet-activating factor is a mediator in tumor necrosis factor/galactosamine-induced lethality.
J. Immunol. 156, 119-123, 1996.
- A.316 VANLANSCHOT, P., BROUHAERT, P., HEIRINCKX, J., KELLERS, R., MIN JOU, W. and PIER, W.
Molecular and immunological characterization of soluble aggregated A/Victoria/3/75 (H3N2) influenza haemagglutinin expressed in insect cells.
Arch. Virol. 141, 1715-1728, 1996.
- A.317 CARRO, M., LOPEZ-SORIANO, F.J., PIER, W. and AGUILES, J.M.
Tumour growth results in changes in placental amino acid transport in the rat: A tumor necrosis factor α mediated effect.
Biochem. J. 313, 77-81, 1996.
- A.318 LEBALLE, P., MOLL, S., JAHN, A., VAN DER MEYDEN, J., TAVENIER, J., PIER, W., DEVOY, S. and TONNEL, A.-D.
EPR-1 is a novel human endothelial cell-specific lectin expressed in lung and regulated by cytokines.
J. Biol. Chem. 271, 20410-20418, 1996.
- A.319 HICK, V., GROSS, A., SIEGHE, S., ROCHETTE, M., PIER, W., DEVOY, S. and DROCK, B.
Abnormal glutathione and sulfate levels after interleukin-6 treatment and in tumor-induced cachexia.
FASEB J. 10, 1219-1226, 1996.
- A.320 VAN CRISTEN, W., BERTHET, R., VAN DE CRAEN, W., VANLANSCHOT, P., BROUHAERT, P., DE VALER, D. and PIER, W.
Functional characterization of the prodomain of interleukin-18-converting enzyme.
J. Biol. Chem. 271, 31515-31522, 1996.
- A.321 RENIEMI, P., FAGGIONI, R., SIROMI, M., MURRAY, C., VANLANSCHOT, P., YAMASHITA, M., SACCO, S., VAN TIL, L.J., N., PIER, W., BROUHAERT, P. and CHILLY, P.
Tumor necrosis factor receptor p55 plays a major role in centrally mediated increases of serum IL-6 and corticosterone after intracerebrum injection of TNF.
J. Immunol. 157, in press, 1996.
- A.322 JOHANSSON, W., WESTERMARK, J., LITTE, A., HALLIN, L., ROIVISTO, L., GELBERG, R., PIER, W., PELTONEN, J., KELLERS, J. and KARL, P.
Collagenase-3 (MMP-13) gene is upregulated by transformed but not by primary human epidermal keratinocytes. Enhancement by tumor necrosis factor- α and transforming growth factor- β .
Submitted.
- A.323 MARCHANT, A., AMARAL, J., CUNHA, C., BRUNY, C., EL NOUE, O., VANLANSCHOT, P., PIER, W., BROUHAERT, P. and GOLDMAN, W.
Betulinic acid: a differentially regulated interferon- γ and tumor necrosis factor production during murine endotoxemia.
Submitted.
- A.324 LIBERT, C., VAN NELLE, W., BROUHAERT, P. and PIER, W.
 α -Antitrypsin inhibits the lethal response to tumor necrosis factor in mice.
J. Immunol. 157, in press, 1996.
- A.325 WERTS, R., BERTHET, R. and PIER, W.
Either the β - or the γ -termini of a loop structure on the highly expressed phage T102 mRNA can contribute to the functional stability of the transcript.
Submitted.
- A.326 WERTS, R., BERTHET, R. and PIER, W.
Optimization of translation initiation in Escherichia coli: the influence of Shine-Dalgarno sequence and upstream ribosomal leak on translational coupling.
Submitted.
- A.327 DECOSTER, E., VANLANSCHOT, P., FLAISANCE, S., VANLANSCHOT, P., BROUHAERT, P. and PIER, W.
Induction of unresponsive T-cells to tumor necrosis factor (TNF) after autocrine TNF expression requires TNF secretion retention.
Submitted.
- A.328 VANLANSCHOT, P., BROUHAERT, P. and PIER, W.
Induced expression of fibronectin intracellularly of the human tumor necrosis factor (TNF) p55 receptor elicits TNF effects in the typhloids of L1210 cells.
Submitted.
- A.329 FLAISANCE, S., VANLANSCHOT, P., BROUHAERT, P., PIER, W. and BROUHAERT, P.
The recombinant signal sequence binding protein (SPB-2) is constitutively bound to the p55 TNF receptor and acts as a negative

J. Biol. Chem. 261, 3345-3348, 1986.

- A.354 RIVIERE, R., FIDIO, V. J., VAN DE CHAUF, M., CORNELIS, S., DECHER, G., LAMM, J. M., GUNDAJAN, A.,
YANDEKCHONG, J. and FIER, W. GUNDAJAN, A.,
Cleavage of F135LH kinase by ICE and CYP11/100-D
during TNF-induced apoptosis.
Submitted.
- A.355 HADAY, M., SATEMI, Z., LAROT, W., FIER, W. and
CONTRENAS, S.
Conversion of the carbohydrate moiety of fungal
glycoproteins to a mammalian type: Evidence for N-
linked GlcNAc-accepting glycans.
Submitted.
- A.356 MEAS, W., DE BRUN, A., SCHMID, J., VERHEIJEN, P.,
CLAYTON, R., FIER, W. and CONTRENAS, S.
Structural characterization of N-linked
oligosaccharides from cellobiohydrolase I secreted by
Trichoderma reesei NRCC 19.
Submitted.
- A.357 BOON, E., VAN LINT, J., GETHMANS, J., ACOSTA, P.,
VANDEKCHONG, J., MEIJER, W., FIER, W.,
VANDEKCHONG, J. and MACHIDA, G.
Identification of a tyrosine phosphorylation site in
the cytoplasmic domain of the human XI 250 tumor
necrosis factor receptor. Phosphorylation by several
members of the src family of protein tyrosine kinases.
Submitted.
- A.358 VAN DE CHAUF, M., VANDEKCHONG, P., DECHER, W., VAN
DE CHAUF, M., VAN LINT, J., MEIJER, W., TROFF, J.,
IMMEL, M., SCHOTTE, P., VAN CHIEKING, W., RIVIERE, R.
and FIER, W.
Characterization of seven murine CED-2/ICE family
members.
Submitted.
- A.359 CAWELS, A., EVERAERT, S., VAN MOLLE, W., BROCKAERT,
P. and FIER, W.
Protection against TNF-induced lethality by methylene
blue, but not when NO synthase is inhibited.
Submitted.
- A.360 CAWELS, A., BROCKAERT, P., SPRINGER, Y., GUTIERREZ-
ZAMOS, J. C. and FIER, W.
Involvement of leukocyte function antigen-1 in systemic
human tumor necrosis factor (TNF) lethality in mice
sensitized to its lethal effects, but not in murine TNF
lethality.
Submitted.

B. SHORT COMMUNICATIONS

- B.1 FIERK, W. and STUCKA, J.
Die neue Polymerease für Ribonucleinsäure und
seine Anwendung bei der Bestimmung von Ribonucleasen.
Naturwiss. 49, 115, 1955.
- B.2 DE BESAGUER, J. and FIERK, W.
Cooperazione des activées phosphatases de
l'Epiderme et des aqueuses phosphatases.
Arch. Biochim. Biophys. 15, 275-276, 1959.
- B.3 FIERK, W. and DE BESAGUER, J.
Nature of the acid phosphatase present in scales of
human skin.
J. Invest. Dermatol. 34, 351, 1960.
- B.4 FIERK, W. and LEPOTTE, L.
The thermal inactivation of bovine ribonucleases.
Arch. Intern. Physiol. Biochim. 49, 282-284, 1961.
- B.5 FIERK, W.
The chromatography of bovine ribonucleases.
J. Chromat. 7, 269-271, 1961.
- B.6 LEPOTTE, L. and FIERK, W.
Séparation des ribonucleases de virus (ribonucleases) par
gel PEG.
Med. Fac. Landbouwwet. 80C 38, 282-284, 1961.
- B.7 VANDERBEEK, P. and FIERK, W.
The distribution of tetracycline present in the
pancreatic ribonuclease digests of bacteriophage M2
RNA and yeast RNA.
Biochim. Biophys. Acta 114, 183-184, 1966.
- B.8 FIERK, W.
Nucleic genetic code of ribonucleic acid
bacteriophage.
Nature 212, 871-873, 1966.
- B.9 DE WACHTER, R., VERHAUSEN, J.P. and FIERK, W.
The 3'-terminal end group of the ribonucleic acid of
the bacteriophage M2.
Arch. Intern. Physiol. Biochim. 76, 176-177, 1968.
- B.10 DE WACHTER, R., VERHAUSEN, J.P. and FIERK, W.
Studies on the bacteriophage M2. V. The 3'-terminal
tetranucleotide sequence of the viral RNA chain.
FEBS Lett. 1, 83-84, 1968.
- B.11 MIN JOU, W., FIERK, W., GOODMAN, H. and SPAIN, P.
Allocation of polypurine tracts to two fragments of
bacteriophage M2 RNA.
J. Mol. Biol. 32, 143-146, 1968.
- B.12 DE BOER, J., GILLIS, G. and FIERK, W.
Q8-virus RNA-afgeknipte RNA-polymerase.
Med. Fac. Landbouwwet. 80C 38, 293-294, 1961.
- B.13 SIEGERS, H. and FIERK, W.
Specific cleavage of ribonucleic acid (RNA) and of viral
RNA in the presence of ribonuclease.
FEBS Lett. 7, 55-56, 1969.
- B.14 MIN JOU, W., CONTRAS, R. and FIERK, W.
The 3'-terminal nucleotide sequence in M2 of
bacteriophage M2 RNA.
FEBS Lett. 9, 221-224, 1970.
- B.15 DE WACHTER, R. and FIERK, W.
The 3'-terminal nucleotide sequence of bacteriophage
M2 RNA.
In "Transcription of Genetic Material", Cold Spring
Harbor Symposium on Quantitative Biology, Vol. 35, Cold
Spring Harbor Laboratory, Cold Spring Harbor, pp. 11-
12, 1970.
- B.16 MIN JOU, W., HARTMAN, C. and FIERK, W.
Studies on the bacteriophage M2. Nucleotide sequence
from the coat protein cistron.
FEBS Lett. 11, 103-107, 1971.
- B.17 DE BOER, J., GILLIS, G. and FIERK, W.
A new method for partial digestion useful for sequence
analysis of polynucleotides.
FEBS Lett. 16, 281-283, 1971.
- B.18 CONTRAS, R. and FIERK, W.
A new method for partial digestion useful for sequence
analysis of polynucleotides.
FEBS Lett. 16, 281-283, 1971.
- B.19 CONTRAS, R., VANDERBEEK, A., MIN JOU, W., DE
WACHTER, R. and FIERK, W.
Studies on the bacteriophage M2. Nucleotide sequence
of a 3'-terminal fragment in M2.
FEBS Lett. 18, 141-144, 1971.
- B.20 SIEGERS, H. and FIERK, W.
Bacteriophage M2 RNA and Escherichia coli 17S
ribosomal RNA have a similar conformation after
reaction with formaldehyde at low pH.
FEBS Lett. 21, 137-140, 1972.
- B.21 DE BOER, J., GILLIS, G. and FIERK, W.
Cellular components involved in the RNA replication
system of bacteriophage M2.
Med. Fac. Landbouwwet. 80C 38, 1021-1027, 1971.
- B.22 CONTRAS, R., VANDERBEEK, A., VOLZGART, G., MIN
JOU, W. and FIERK, W.
Studies on the bacteriophage M2. Some nucleotide
sequences from the RNA-polymerase gene.
FEBS Lett. 24, 119-121, 1972.
- B.23 VANDERBEEK, A. and FIERK, W.
Further evidence on the role of the A protein in
bacteriophage M2 replication.
FEBS Lett. 30, 89-92, 1972.

- 0.24 DE BOEVER, J., VAN EMDELD, J., GILLIS, E. and PIERIS, W.
The affinity of host factor for 18 viral RNA.
Med. Fac. Landbouwwet. RUC 28, 1873-1879, 1973.
- 0.25 PIERIS, W.
The choice of virus source in M31 RNA.
Matur. Rev. Sint. 24, 186, 1973.
- 0.26 VOLCKAERT, C. and PIERIS, W.
Studies on the bacteriophage M31. G-U-C as the initiation codon of the A-protein cistron.
Virus Lett. 15, 81-86, 1973.
- 0.27 VAN EMDELD, J., GILLIS, E. and PIERIS, W.
Template related effects in the replication reaction of Q β -virus RNA.
Med. Fac. Landbouwwet. Gent 18, 1171-1176, 1974.
- 0.28 NIM JON, W. and PIERIS, W.
Sequence determination of Q β -rich oligonucleotides by means of the Rothwell modification.
Virus Lett. 66, 77-81, 1974.
- 0.29 VAN EMDELD, J., WERREICHAERT, J., DEVOS, M., GILLIS, E. and PIERIS, W.
A new method for sequence analysis of plant virus RNA.
Med. Fac. Landbouwwet. RUC 22, 1193-1197, 1974.
- 0.30 HADJIANAN, C. and PIERIS, W.
Evidence for splicing of SV40 165 mRNA.
Matur. 211, 70-73, 1976.
- 0.31 PIERIS, W. and CROSLAND, H.
On codon usage.
Matur. 211, 328, 1976.
- 0.32 CONTRERAS, R., COLE, C., BIER, P. and PIERIS, W.
Nucleotide sequence analysis of two Sialian virus 40 mutants with deletions in the late region of the genome.
J. Virol. 28, 783-793, 1978.
- 0.33 VAN MEUVENSSEM, M., COLE, C., BIER, P. and PIERIS, W.
Nucleotide sequence analysis of two Sialian virus 40 mutants with deletions in the region coding for the carboxyl terminal of the T antigen.
J. Virol. 30, 838-841, 1979.
- 0.34 VAN EMDELD, J., WERREICHAERT, M., DEVOS, M. and PIERIS, W.
The primary structure of the satellite tobacco necrosis virus genome.
Med. Fac. Landbouwwet. RUC 44, 801-810, 1979.
- 0.35 CONTRERAS, R., CHENOUTE, M. and PIERIS, W.
A simple apparatus for injection of neonatal quail-like trachea into *Isoparus laevis* oocytes.
Anal. Biochem. 122, 183-187, 1981.
- 0.36 GUERIN, D. and PIERIS, W.
Expression and secretion of human fibroblast 81 interaction in monkey cells after transfection with a recombinant SV40 plasmid vector.
J. Mol. Appl. Genet. 1, 183-188, 1981.
- 0.37 MULDER, P., VAN EMDELD, J. and PIERIS, W.
The sequence specificity of endonuclease Cof and CofII isolated from *Chloroflexus aurantiacus*.
Gene 18, 91-96, 1981.
- 0.38 CORTEOT, J., DE MIR, L. and PIERIS, W.
In vitro correlations of processing of human gamma-interferon 3 enhancer to biological activity.
Virology 123, 466-470, 1983.
- 0.39 VAN ROY, P., FRANK, L. and PIERIS, W.
Metabolic turnover of phosphorilation sites in Sialian virus 40 large T antigen.
J. Virol. 45, 443-448, 1983.
- 0.40 TAVENIER, J., CHREYSEN, D., HUEBNER, P., VAN DER MEULEN, J. and PIERIS, W.
Deletion mapping of the inducible promoter of human Matur. 301, 414-416, 1983.
- 0.41 ROBER, J.-J., COLLINS, T., GIBSON, M. A., COCHRAN, R. J., GILLIS, J. D., PIERIS, W., CLAYTON, C., FRANK, A. M., BURROFF, S. J. and REISS, E. J.
Lymphocyte recognize human vascular endothelial and dermal fibroblast as antigen induced by recombinant immune interferon.
Matur. 305, 718-720, 1983.
- 0.42 AMELIOT, P., VAN EMDELD, J. and PIERIS, W.
SV-4, a new satellite tobacco necrosis virus variant.
Med. Fac. Landbouwwet. RUC 48, 101-109, 1983.
- 0.43 DEVOS, M., PLATTINCK, C. and PIERIS, W.
Induction of cytopathic cells by pure recombinant human interferon 3.
Eur. J. Immunol. 14, 1059-1060, 1984.
- 0.44 TAVENIER, J. and PIERIS, W.
The presence of homologous regions between interferon sequences.
Carlsberg Res. Commun. 43, 159-164, 1984.
- 0.45 CONTRERAS, R. and PIERIS, W.
A method for fast and accurate RNA solution from agarose gels by centrifugal filtration.
Bio/Technology 1, 1016-1018, 1985.
- 0.46 AMELIOT, P., VAN EMDELD, J. and PIERIS, W.
Synthesis of satellite tobacco necrosis virus (STNV) RNA-like transcripts in *Escherichia coli*.
Med. Fac. Landbouwwet. RUC 50, 1381-1383, 1984.

- B.43 DEBRAYE, M., SIMONS, G., DEVOS, R., PLATTINCK, G., REHAUT, R., TAVERNIER, J., and FIEBS, W.
Cloning and structure of a mouse Interleukin-2 Chromosomal gene.
Mol. Biol. Sup. 12, 59-61, 1984.
- B.48 MULLER, R., KASHIMOTO, A. and FIEBS, W.
Synthesis and maturation of recombinant human tumor necrosis factor in eukaryotic systems.
FEBS Lett. 137, 99-101, 1984.
- B.49 SMITH, D.A., DISTER, M., FIEBS, W. and FAGLIOMI, C.
Species specificity of human and murine tumor necrosis factor. A comparative study of tumor necrosis factor receptors.
J. Biol. Chem. 259, 11811-11816, 1984.
- B.50 COOPERMAN, E., MARTEL, M., VAN ROY, P., GAO, J. and FIEBS, W.
Invasive and metastatic capability of rat cells before and after transfection with bovine papilloma virus 1 DNA.
Arch. Biol. 97, 381, 1988.
- B.51 MALKOWSKY, M., LOVELAND, B., MORTIN, M., ANDERSON, G.L., GAO, J., GAO, P. and FIEBS, W.
Recombinant Interleukin-2 directly augments the cytotoxicity of human monocytes.
Nature 323, 362-365, 1987.
- B.52 PLATTINCK, G., DELLECO, M., TAVERNIER, J., MARHOLZ, M. and FIEBS, W.
Recombinant tumor necrosis factor can induce Interleukin 2 receptor expression and cytotoxic activity in a rat x mouse T cell hybrid.
Eur. J. Immunol. 17, 1835-1838, 1987.
- B.53 SUTYI, P., BESANT, R., VAN ROY, P. and FIEBS, W.
Reduced tumor necrosis factor-induced cytotoxicity by inhibitors of the arachidonic acid metabolism.
Biochem. Biophys. Res. Commun. 149, 718-723, 1987.
- B.54 HETZEL, A., BOTTENBERG, J.W., FIEBS, W. and DUMONT, J.E.
Modulation of tumor necrosis factor cytotoxicity in L937 cells by bacterial toxins, hydrocortisone and inhibitors of arachidonic acid metabolism.
Biochem. Biophys. Res. Commun. 149, 615-622, 1987.
- B.55 DEVOS, R., TAVERNIER, J. and FIEBS, W.
Release of DNA polymerase I during synthesis of dsDNA.
Mol. Acids Res. 16, 1838, 1988.
- B.56 EUFFE, P., VAN ROY, P. and FIEBS, W.
Tumor necrosis factor and interleukin 2 activate phospholipase in rat chondrocytes.
FEBS Lett. 212, 24-28, 1988.
- B.57 VANDERKILDE, P., JAPANEK, B., DEVOS, R., BLAU, R. and FIEBS, W.
Interleukin 2 acts as an autocrine growth factor for myeloma cells. An Epstein-Barr virus-transformed human B cell line.
Eur. J. Immunol. 18, 1027-1031, 1988.
- B.58 VAN OSTADE, X., TAVERNIER, J. and FIEBS, W.
Two converted tryptophan residues of tumor necrosis factor and receptor are not involved in the biological activity.
FEBS Lett. 238, 347-352, 1988.
- B.59 HETZEL, A., DUMONT, J.E., BOTTENBERG, J.W., FIEBS, W. and DUMONT, J.E.
Pepin phosphorylation of a 22 kDa protein induced by tumor necrosis factor.
FEBS Lett. 237, 173-176, 1988.
- B.60 KAMMAY, M., DE SUT, W., FIEBS, W. and HERNAN, A.C.
Inflammation, chemotaxis, and tumor necrosis factor in rabbit skin in vivo.
J. Exp. Med. 169, 223-228, 1989.
- B.61 BROUCHART, P., SPRINGS, D.B., DUMONT, J.E., KUFF, D.W. and FIEBS, W.
Circulating interleukin 2 during a continuous infusion of tumor necrosis factor and interferon γ .
J. Exp. Med. 169, 229-232, 1989.
- B.62 BROUCHART, P., C. FIEBS, W., EVERAERT, D., TAVERNIER, J. and FIEBS, W.
A role for interleukin 1 in the in vivo actions of tumor necrosis factor.
Lymphokine Res. 8, 349-356, 1989.
- B.63 EVERAERT, D., BROUCHART, P., SHUK, A. and FIEBS, W.
Four different interleukin-1 species, consisting of the lethal action of tumor necrosis factor.
Biochem. Biophys. Res. Commun. 163, 330-335, 1989.
- B.64 ROBBE, R., HETZEL, A., LECHE, R., FIEBS, W., BOTTENBERG, J.W. and DUMONT, J.E.
Tumor necrosis factor induces the phosphorylation of the stress protein in endothelial cells: Possible role in protection against cytotoxicity.
Biochem. Biophys. Res. Commun. 161, 301-308, 1989.
- B.65 LECHE, R., VANDERKILDE, P., FLAMMAR, V., ROBBE, R., LECHE, R., ANDERSON, G., TAVERNIER, J., FIEBS, W. and COOPERMAN, E.
Hypothermia and hypoxemia induced by anti-CD1 monoclonal antibody in mice: Role of tumor necrosis factor.
Eur. J. Immunol. 18, 107-110, 1988.
- B.66 VANDERKILDE, P., CHACOE JR, E.J., ROBBE, R., SUTYI, P., VAN ROY, P. and FIEBS, W.
Cytotoxic activity of tumor necrosis factor is inhibited by antisense derivative antisense mRNA.

- 8.67 BEVAERT, R., SUFFYS, P., VAN ROY, F. and FIERZ, W.
Inhibition by glucocorticoids of tumor necrosis factor-mediated cytotoxicity. Evidence against lipocortin involvement.
FEBS Lett. 181, 91-94, 1990.
- 8.68 LIBERT, C., BROUHAERT, P., SHAW, A. and FIERZ, W.
Induction of interleukin 6 by human and murine recombinant interleukin 1 in mice.
Eur. J. Immunol. 20, 691-694, 1990.
- 8.69 PARANT, P., TAVENIER, J., FIERZ, W. and PARANT, R.
Comparative activity of human and murine tumor necrosis factor in toxicity and anti-infectious assays in mice.
Microb. Pathogen. 8, 181-188, 1990.
- 8.70 GONZA, J., SIBRA, R., FIERZ, W., CAULDIE, J. and KOJ, A.
Transforming growth factor- α and epidermal growth factor modulate basal and interleukin-6-induced cathecol uptake and acute phase protein synthesis in cultured rat hepatocytes.
FEBS Lett. 244, 49-50, 1990.
- 8.71 VANDENHART, P., GUISSE, Y., DESCIENCO, W., BAUM, G., VANDEKROON, J. and FIERZ, W.
Responses of murine cell lines to an IL-1/IL-2-induced factor in a rat/mouse hybridoma (PCHO) differential induction of cytokines by human IL-1 α and IL-1 β and partial amino acid sequence of rat GM-CSF.
Lymphokine Res. 7, 381-389, 1990.
- 8.72 COLMAN, M., VANDENHART, P., MULLART, J., AMAROU, E., VANDEKROON, J., MORTIER, J., VANDEKROON, J.L., FIERZ, W. and STIZOL, Y.
Interleukin-6 secretion of interleukin-4 during continuous subcutaneous perfused dialysis.
Nephron 56, 376-380, 1990.
- 8.73 DEVOS, R., TAVENIER, J., PLATTINCK, G., VAN DER HEYDEN, J., BOLINK, A. and FIERZ, W.
Expression of the murine interleukin-3 receptor on xenopus laevis oocytes.
Biochem. Biophys. Res. Commun. 171, 570-575, 1990.
- 8.74 CARLO-STELLA, C., MANCINI, L., ALBERTI, C., FRASSON, F., FIERZ, W. and STIZOL, Y.
Growth of COH1 acute myeloblastic leukemia colony-forming cells in response to recombinant hematopoietic growth factors.
Leukemia 4, 561-568, 1990.
- 8.75 HOROWITZ, T., KORTA, M., SOJ, A., FIERZ, W., CAULDIE, J. and BAUM, G.
Effects of interleukin-6 and leukemia inhibitory factor on the acute phase response and DNA synthesis in cultured rat hepatocytes.
Lymphokine Cytokine Res. 10, 23-28, 1991.
- 8.76 DEVOS, R., VANDENHART, P., BOLINK, A., PLATTINCK, G., VAN DER HEYDEN, J., FIERZ, W. and MULLART, J.
Acute phase response of a mouse interleukin-3 receptor protein.
Eur. J. Immunol. 21, 1115-1117, 1991.
- 8.77 VANDENHART, P. and FIERZ, W.
Is cytokine-mediated during Alzheimer's disease due to an IL-1/IL-6-mediated acute phase response in the brain? Today 12, 217-219, 1991.
- 8.78 VANDENHART, P., MULLART, G. and FIERZ, W.
Tumor necrosis factor-induced interleukin-6 expression and cytotoxicity follow a common signal transduction pathway in L1210 cells.
Biochem. Biophys. Res. Commun. 170, 981-1000, 1991.
- 8.79 BROUHAERT, P., LIBERT, C., VANDEKROON, J. and FIERZ, W.
Selective species specificity of tumor necrosis factor for toxicity in the mouse.
Lymphokine Cytokine Res. 11, 191-196, 1992.
- 8.80 BIRNEY, R., SCHULZ-OSTHOFF, R., VAN ROY, F. and FIERZ, W.
Synthetic induction of interleukin-6 by tumor necrosis factor and interleukin-1 α in alveolar cells in the triggering and exacerbation of psoriasis by lithium treatment.
Eur. J. Immunol. 22, 3181-3184, 1992.
- 8.81 VANDENHART, P., MULLART, G. and FIERZ, W.
TNF-mediated IL-6 gene expression and cytotoxicity are co-inducible in TNF-resistant L1210 cells.
FEBS Lett. 302, 335-338, 1992.
- 8.82 BROUHAERT, P., G., FIERZ, W. and LEBJUNE, F.J.
Colony vaccine and TNF therapy.
Nature 358, 610, 1992.
- 8.83 VANDEKROON, J., BARTENS, G., MIN JOU, W. and FIERZ, W.
Recombinant secreted hematopoietin protects mice against a lethal challenge of influenza virus.
Vaccine 11, 1183-1187, 1993.
- 8.84 GUISSE, Y., LIBERT, C., MULLART, G., SCHULZ-OSTHOFF, R., VAN ROY, F., VAN DER HEYDEN, J., PLATTINCK, G., CARLO-STELLA, C., TAVENIER, J., FIERZ, W., DEVOS, R. and FIERZ, W.
Expression, purification and cytotoxicity of fully active, glycosylated human interleukin-6.
FEBS Lett. 331, 49-51, 1993.
- 8.85 TIE-HE, L., ROCCCH, E., MIN JOU, W. and FIERZ, W.
Expression of neuraminidase of influenza virus in insect cells.
Chin. J. Biochem. Biophys. 15, 337-341, 1993.

- 8.00 STEIDLER, L., WELLS, J. M., PARSHAKUR, A.,
VANDERKAMME, J., FIERS, W. and REMAUT, E.
Secretion of biologically active serum intercalin-3
by Lactococcus lactis subsp. lactis.
Appl. Environ. Microbiol. 41, 1417-1419, 1985.
- 8.07 DE VALLEE, D., REINING, K., VAN OSTERINGE, W.,
CONTRETT, S., BEYRENT, B. and FIERS, W.
338, An inhibitor of cell death, self-associated by
its zinc finger domain.
FEBS Lett. 186, 31-34, 1985.
- 8.08 STEIDLER, L., VO, M., FIERS, W. and REMAUT, E.
The expression of the photolabile luciferase gene
in Staphylococcus aureus Cowan I allows the
development of a live explantable tool for
immunodetection.
Appl. Environ. Microbiol. 41, 1354-1358, 1985.
- 8.09 CLAUS, M., GRILL, M., FANCKAER, C., FIERS, W.,
SCHURIM, P. and RISKU, W.
Synergistic induction of endothelial tissue factor by
tumor necrosis factor and vascular endothelial growth
factor: functional analysis of the tumor necrosis
factor receptor.
FEBS Lett. 184, 334-338, 1985.

C. a. BOOKS (CONTINUED CHAPTERS, EDITORIALS)

- C. a. 1 FIELDS, W.
The structure of viral RNA.
In Ouchetto, J. (ed.), "Structural Studies on Nucleic Acids and Other Biopolymers", Physical-Chemical Properties of nucleic acids, Vol. 1, Academic Press, London-New York, pp. 111-131, 1973.
- C. a. 2 FIELDS, W.
RNA bacteriophages.
In King, R.C. (ed.), "Handbook of Genetics", Vol. 1, Plenum Press, Inc., New York, pp. 171-196, 1974.
- C. a. 3 FIELDS, W.
Chemical structure and biological activity of bacteriophage MS2 RNA.
In Lindner, W. (ed.), "RNA Phages", Cold Spring Harbor Laboratory, Cold Spring Harbor, pp. 151-184, 1975.
- C. a. 4 FIELDS, W.
Nucleic acid sequences.
In Felsen, G.D. (ed.), "Nucleic Acids", Handbook of Biochemistry and Molecular Biology, Vol. 3, CRC Press, Cleveland, pp. 314-353, 1974.
- C. a. 5 FIELDS, W. and FIELDS, W.
Structure and function of viral nucleic acids.
In Clark, R.F.C. (ed.), "Biochemistry of Nucleic Acids", International Review of Biochemistry, Vol. 17, II, University Park Press, Baltimore, pp. 12-131, 1978.
- C. a. 6 FIELDS, W.
Structure and function of RNA bacteriophages.
In Fraenkel-Conrat, M. and Wagner, R.R. (eds.), "Comprehensive Virology", Vol. 13, Plenum Publishing Corporation, New York, pp. 89-204, 1979.
- C. a. 7 DE WINTER, R. and FIELDS, W.
Two-dimensional gel electrophoresis of nucleic acids.
In Rickwood, D. and Hames, B.D. (eds.), "Gel Electrophoresis of Nucleic Acids: A Practical Approach", IRL Press Limited, Oxford-Washington, pp. 77-116, 1981.
- C. a. 8 REMAUT, E., STANISSEY, F., SIMONS, G. and FIELDS, W.
Use of the phage lambda P1 promoter for high-level expression of human interferon in Escherichia coli.
In Festschrift, 3, (ed.), "Interferon", Methods in Enzymology, Vol. 115, Part C, Academic Press, Inc., Orlando, pp. 366-375, 1986.

- C. a. 9 TAYLOR, J., FRANKS, R., L. HAMMING, A., VAN DEN HERTUM, J., MULLEN, R., ROYSHAM, R., VAN VLIET, A., BAUDER, R. and FIELDS, W.
Isolation and expression of the genes coding for mouse and human tumor necrosis factor (TNF) and biological properties of recombinant TNF.
In Gubb, D. and Gaudel, D.V. (eds.), "Molecular Cloning and Analysis of Tumor Necrosis Factor", Academic Press, Inc., Orlando, pp. 161-196, 1987.
- C. a. 10 REMAUT, E., WERNER, A. and FIELDS, W.
Expression of heterologous unfused protein in Escherichia coli.
In W. B. and Gaudel, D. (eds.), "Recombinant DNA Part B: Methods in Enzymology", Vol. 153, Academic Press, Inc., San Diego, pp. 118-131, 1987.
- C. a. 11 BALKWILL, F.R. and FIELDS, W.
[unintelligible] Biological Response Modifiers.
In "Cancer Surveys", Imperial Cancer Research Fund, London, Vol. 6 (1), pp. 111-127, 1989.
- C. a. 12 CONTRAS, M., DONDOLIN, J. and FIELDS, W.
Cloning and expression of cytokine genes.
In Balkwill, F.R. (ed.), "Cytokines: A Practical Approach", IRL Press, Oxford, pp. 1-16, 1991.
- C. a. 13 FIELDS, W.
Protein structure and structure-function analysis of TNF and lymphotxin.
In Aggarwal, B.B. and Vilcek, J. (eds.), "Tumor Necrosis Factor: Structure, Function, and Mechanism of Action", Marcel Dekker, Inc., New York/Basel/Stuttgart, pp. 39-93, 1991.
- C. a. 14 FIELDS, W.
Tumor Necrosis Factor.
In Sica, E. (ed.), "The Natural Immune System: Humoral Factors", IRL Press, Oxford, pp. 83-119, 1991.
- C. a. 15 FIELDS, W. and GURRIGAN, M.A.
[eds.] "Tumor Necrosis Factor: Molecular and Cellular Biology and Clinical Potency".
Proceedings of the 1th International Conference on Tumor Necrosis Factor, Veldhoven (The Netherlands), May 1-6, 1991. Harger, Basel, 356 pp., 1991.
- C. a. 16 FIELDS, W.
Biologic therapy with TNF: Practical studies.
In DeVita, D.T., Hellman, S. and Rosenberg, S.A. (eds.), "Biologic Therapy of Cancer", 2nd ed., J.B. Lippincott, Philadelphia, pp. 193-217, 1993.
- C. a. 17 CONTRAS, M., DEMOJON, J. and FIELDS, W.
Cloning and expression of cytokine genes.
In Balkwill, F.R. (ed.), "Cytokines: A Practical Approach", 2nd ed., IRL Press, Oxford, pp. 1-18, 1991.

C. B. REVIEWS CONCERNING OWN RESEARCH

- C.B.1 PIERIS, W.
De ribonucleïnezuur-afbrekende enzymen van gerst.
Proefschrift Agr. NO. 305, 284 blz. 1940.
- C.B.2 PIERIS, W.
Studia over het verband tussen chemische structuur
en biologische activiteit van het nucleïnezuur van
de bacteriophage 52-118.
Natuurwet. Tijdschrift 15, 3-116, 1943.
- C.B.3 PIERIS, W., DE WACHTER, R., MIN JOU, W. and VAN
STIVINGALE, B.
The structure of bacteriophage RNA.
1st International Congress for Virology (Amsterdam),
in Heinich, J. L. (ed.), "International Virology",
Vol. 1, Springer, 1961, pp. 30-35, 1961.
- C.B.4 PIERIS, W., VAN MONTAGU, M., DE WACHTER, R., HAZENMAN,
G., MIN JOU, W., HESSEN, E., REYAUT, E.,
VANDERBEEK, A. and VAN STIVINGALE, B.
Studies on the primary structure and the replication
mechanism of bacteriophage RNA.
In "The Mechanism of Protein Synthesis", Cold Spring
Harbor Symposium on Quantitative Biology, Vol. 16,
Cold Spring Harbor Laboratory, Cold Spring Harbor,
pp. 433-453, 1951.
- C.B.5 PIERIS, W., CONTRERAS, R., DE WACHTER, R., HAZENMAN,
G., HERRIGERT, J., MIN JOU, W. and VANDERBEEK, A.
Recent progress in the sequence determination of
bacteriophage MS2 RNA.
Biochimica 33, 293-304, 1971.
- C.B.6 PIERIS, W., CONTRERAS, R., DE WACHTER, R., HAZENMAN,
G., HERRIGERT, J., MIN JOU, W., VANDERBEEK, A.
Studies on the structure of bacteriophage MS2 RNA,
1st International Congress for Virology, Amsterdam
(Amstert),
in Heinich, J. L. (ed.), "International Virology 2",
Springer, 1961, pp. 1-11, 1971.
- C.B.7 MIN JOU, W. and PIERIS, W.
The complete nucleotide sequence of a bacteriophage
gene.
In "Encyclopedia della scienza e della Technica",
1971.
- C.B.8 PIERIS, W., CONTRERAS, R., DE WACHTER, R., HAZENMAN,
G., HERRIGERT, J., MIN JOU, W., VANDERBEEK, A.,
VOLTERRA, G. and HERRIGERT, R.
Structure and function of the bacteriophage MS2 RNA.
in Heinich, J. L., Ochoa, J. and Ore, J. (eds.),
"Viral Replication and Cancer", Editorial Labor,
Barcelona, pp. 11-22, 1973.

- C.B.9 PIERIS, W., DANDA, R., ROGIER, R., VAN DE VOORDE, A.,
VAN HERRIGERT, J., VAN HERRIGERT, W., VOLCKERT,
G. and VANDERBEEK, A.
Approaches to the sequence determination of MS2
RNA.
In "Tumor Viruses", Cold Spring Harbor Symposium on
Quantitative Biology, Vol. 32, Part 1, Cold Spring
Harbor Laboratory, Cold Spring Harbor, pp. 119-124,
1971.
- C.B.10 CRAMON, R. L., AGOSTINI, R. J., MULLER, C., HERRIGERT,
G. L., VANDERBEEK, A., DE WACHTER, R., PIERIS, W.
Studies on in vitro transfection by DNA and RNA
fragments of human adenovirus and Simian Virus 40.
In "Tumor Viruses", Cold Spring Harbor Symposium on
Quantitative Biology, Vol. 32, Part 1, Cold Spring
Harbor Laboratory, Cold Spring Harbor, pp. 417-420,
1971.
- C.B.11 PIERIS, W.
Bacteriophage MS2: Een modelstelsel voor de studie
van de structurele relatie van het officiële
zetaarrest.
Chem. Weekblad 30, VI-VS, 1974.
- C.B.12 PIERIS, W., ROGIER, R., SOEDER, E., VAN DE VOORDE,
A., VAN HERRIGERT, W., VAN HERRIGERT, J.,
VOLCKERT, G. and VANDERBEEK, A.
Nucleotide sequence analysis of MS2 RNA.
10th FEBS Meeting, Paris (France), July 30-31, 1973.
In "Preconferences", pp. 13-13, 1973.
- C.B.13 PIERIS, W.
Recent developments in MS2-RNA: Die
vollständige Struktur eines Virus-Genoms und ihre
biologische Bedeutung.
Bechr. Chem. Techn. 36, 311-316, 1974.
- C.B.14 PIERIS, W.
Die Formel eines Virus: Die RNA des Bacteriophagen
MS2 - die vollständige chemische Struktur eines
Virus-Genoms und ihre biologische Bedeutung.
Medizin in unserer Zeit 1, 19-21, 1972.
- C.B.15 VAN DE VOORDE, A., CONTRERAS, R., HAZENMAN, G.,
ROGIER, R., VAN HERRIGERT, W., VAN HERRIGERT, J.,
VOLCKERT, G., VANDERBEEK, A. and PIERIS, W.
Structural organization of the MS2 genome.
1st Colloquium de l'Institut National de la Santé et
de la Recherche Médicale, Gignoux (France), July 9,
1972.
In May, P., Monier, R. and Wolf, R. (eds.), "Early
Protein of Oncogenic DNA Viruses", INSERM, Vol. 49,
INSERM, Paris, pp. 17-30, 1972.

- C.b.16
 CONTRERAS, R., KATZMAN, C., ROGLIER, A., VAN DE
 VOORSE, A., VAN HERTEN, M., VAN HERTEN, J.,
 VOLKERT, G., VERBEEK, H. and PERS, M.
 Complete nucleotide sequence of SV40 DNA.
 12th FEMS Meeting, Dresden (Germany), July 2-7,
 1981.
 In Rosenthal, A. (ed.), "Gene Function", Federation
 of European Biochemical Societies, Vol. 31, Pergamon
 Press, Oxford-New York, pp. 119-125, 1981.
- C.b.17
 MIN JOU, M., VERHOEVEN, M., DEVOS, R., SMAN, E.,
 KATZMAN, C., VAN HERTEN, M., PANG, R. and PERS, M.
 Cloning and DNA nucleotide sequence analysis of the
 haemagglutinin and neuraminidase genes of Influenza A
 strains.
 International Workshop on Structure and Variation in
 Influenza Virus, Thredbo (Australia), December 18-
 23, 1980.
 In Iavar, G. and Alr, C. (eds.), "Structure and
 Variation in Influenza Virus", Elsevier/North-
 Holland, New York, pp. 61-89, 1980.
- C.b.18
 MIN JOU, M., VERHOEVEN, M., PANG, R., DEVOS, R.,
 KATZMAN, C., SMAN, E. and PERS, M.
 Complete primary structure of the haemagglutinin from
 two different strains of Influenza A type virus.
 In "Protides of the Biological Fluids", Pergamon
 Press, New York, pp. 141-147, 1980.
- C.b.19
 MIN JOU, M., VERHOEVEN, M., PANG, R. and PERS, M.
 Drift and shift of Influenza A-type virus
 with single nucleotide substitution in the HA gene.
 12th FEMS Meeting, Dresden (Germany), July 2-7,
 1981.
 In Rosenthal, A. (ed.), "Gene Function", Federation
 of European Biochemical Societies, Vol. 31, Pergamon
 Press, Oxford-New York, pp. 119-125, 1981.
- C.b.20
 DEBRAS, R., DEVOS, R., REMAUT, E., SMAN, E.,
 KATZMAN, C., VERHOEVEN, M., PANG, R., DEVOS, R.,
 VOLKERT, G., VERBEEK, H. and PERS, M.
 Isolation and characterization of a human fibroblast
 interferon gene and its expression in Escherichia
 coli.
 Res. Infect. Dis. 3, 1164-1173, 1981.
- C.b.21
 MIN JOU, M., VERHOEVEN, M., PANG, R., DEVOS, R.,
 KATZMAN, C. and PERS, M.
 Shift and drift in Influenza A virus.
 12th Symposium of the Society for General
 Microbiology, Edinburgh (United Kingdom), September
 1981.
 In Corti, M.J., Collins, J.F. and Roseley, S.E. (eds.),
 "Molecular and Cellular Aspects of Microbial
 Evolution", Cambridge University Press, Cambridge,
 pp. 103-111, 1981.
- C.b.22
 PERS, M., PANG, R., MIN JOU, M., KATZMAN, C.,
 VERHOEVEN, M., DEVOS, R. and VAN HERTEN, J.
 Drift and shift of Influenza A virus at the genome
 level.
 In Iavar, G. and Alr, C. (eds.), "Structure and
 Variation in Influenza Virus", Elsevier/North-
 Holland, New York, pp. 119-125, 1981.
- C.b.23
 DEBRAS, R., REMAUT, E., KATZMAN, C., VERHOEVEN, M.,
 PANG, R., DEVOS, R., VOLKERT, G., VERBEEK, H. and
 PERS, M.
 Isolation and characterization of a human fibroblast
 interferon gene and its expression in Escherichia
 coli.
 Res. Infect. Dis. 3, 1164-1173, 1981.
- C.b.24
 REMAUT, E., DEBRAS, R., KATZMAN, C., VERHOEVEN, M.,
 PANG, R., DEVOS, R., VOLKERT, G., VERBEEK, H. and
 PERS, M.
 Isolation and characterization of a human fibroblast
 interferon gene and its expression in Escherichia
 coli.
 Res. Infect. Dis. 3, 1164-1173, 1981.

- C.B.27 PIERIS, M., DEVOS, R., CHEROUTRE, M., DEGRAVE, M., DUEZINK, F., CHRYSEN, G., FLATTING, G., REMOUT, E., SCAMILL, S., SIMONS, G., STAMMERS, F., TAVENIER, J., and VAN DER WOUDE, J. Molecular-biological studies on human fibroblast interferon, immune interferon and interferon 3 genes. In De Maeyer, E. and Schellkens, H. (eds.), "The Biology of the Interferon System 1983", Elsevier Science Publishers, Amsterdam, pp. 3-6, 1983.
- C.B.28 COXNET, J., DE WIT, L., TAVENIER, J., and PIERIS, M. Human fibroblast interferon mRNA in poliovirus-induced cells. In De Maeyer, E. and Schellkens, H. (eds.), "The Biology of the Interferon System 1983", Elsevier Science Publishers, Amsterdam, pp. 11-24, 1983.
- C.B.29 CHEROUTRE, M., DEVOS, R., FLATTING, G., SCAMILL, S., DEGRAVE, M., TAVENIER, J., YATA, Y., and PIERIS, M. Molecular cloning of human lymphokines. In Mowbray and Rosenburg (eds.), "Gene Expression", Vol. VII, Alan R. Liss, Inc., New York, pp. 417-444, 1983.
- C.B.30 VAN BOMPUY, L., VAN LOU, W., VERHOEF, M., KUIJZER, D., and PIERIS, M. Molecular variation of influenza surface antigens. *Virus* 9, 414-417, 1981.
- C.B.31 VAN BOMPUY, L., VAN LOU, W., VERHOEF, M., KUIJZER, D., and PIERIS, M. Control of expression of plant virus-coded information in E. coli. *Mol. Cell. Biochem.* 88, 141-153, 1981.
- C.B.32 PIERIS, M., VAN BOMPUY, L., and VAN LOU, W. And viruses as vectors in genetic engineering of plants. In "Research and Training Programme in Biotechnology Engineering", Progress Report, Vol. 1: Research, Commission of the European Communities, Brussels, pp. 134-138, 1983.
- C.B.33 PIERIS, M., REMOUT, E., DEVOS, R., STAMMERS, F., SIMONS, G., SCAMILL, S., CHEROUTRE, M., FLATTING, G., TAVENIER, J., DEGRAVE, M., DUEZINK, F., and VAN DER WOUDE, J. High level synthesis of proteins in heterologous cells using genetic engineering approaches. In Grosswald, M. (ed.), "Molecular Chemistry for the Future", Pergamon Press, Oxford-New York, pp. 311-341, 1984.
- C.B.34 VAN DE VOORDE, A., DE WIT, L., DE CROOTE, G., POLLET, D., DE BRON, M., and PIERIS, M. Monoclonal antibodies as a tool in the detection of human fibroblast alkaline phosphatase. In Fishbein, S. W. (ed.), "Progress in Clinical and Biological Research: Essays on Human Alkaline Phosphatase", Alan R. Liss, Inc., New York, pp. 199-207, 1984.
- C.B.35 VAN DE VOORDE, A., DE WIT, L., DE CROOTE, G., POLLET, D., DE BRON, M., DE BOEVER, J., VANDEKAPPELE, D., and PIERIS, M. The application of a monoclonal antibody for the detection of human fibroblast alkaline phosphatase in cells and tumor extracts of cancer patients. In Fishbein, S. W. (ed.), "Progress in Clinical and Biological Research: Essays on Human Alkaline Phosphatase", Alan R. Liss, Inc., New York, pp. 29-38, 1984.
- C.B.36 PIERIS, M., DEVOS, R., CHEROUTRE, M., DEGRAVE, M., FLATTING, G., REMOUT, E., SCAMILL, S., SIMONS, G., and TAVENIER, J. Molecular biology of the human interferon 3 and interferon 2 genes. In Feilicoff, E., Fildes, W. H., and Cillo, S. (eds.), "Lymphokines", Tokyo: Sando Research Center, Vol. 49, Institut Scientifique Roussel, Paris, pp. 11-19, 1984.
- C.B.37 PIERIS, M. Cell differentiation and gene expression: a brief overview. In Conference on Differentiation and Disease. Gent (Belgium), November 28 - December 1, 1984. In Verbruggen, G. and Vervo, K. M. (eds.), "Regenerative Toxins", Vol. 3, Elsevier Science Publishers, Amsterdam, pp. 3-7, 1984.
- C.B.38 PIERIS, M., DEVOS, R., CHEROUTRE, M., DEGRAVE, M., GUYEN, D., FLATTING, G., REMOUT, E., SCAMILL, S., SIMONS, G., STAMMERS, F., TAVENIER, J., and VAN DER WOUDE, J. Cloning, expression and characterization of human interferon-beta, interferon-gamma and interferon-2 genes. In Battistoni, U., Bittum, F. J., and Chlone, R. (eds.), "Contributions of Modern Biology to Medicine. New Approaches in Pathogenesis, Diagnosis and Therapy in Immunological and Neoplastic Diseases", Sando Research Publications, Vol. 17, Raven Press, New York, pp. 21-23, 1983.
- C.B.39 PIERIS, M., DEVOS, R., REMOUT, E., STAMMERS, F., TAVENIER, J., MARINHO, A., GUILLI, Y., and VAN DER WOUDE, J. Interferon genes and its synergism with tumor necrosis factor in human endocrine systems. In Duenkel, F. and Giesel, G. B. (eds.), "The Interferon System", Sando Research Publications, Vol. 14, Raven Press, New York, pp. 143-151, 1984.

- C.B.40 FIEBS, W., DEGRAVE, W., DEVOS, R., FLATTECK, C., CHENOMUTH, W., TAVENIER, J., SIMONE, G., and REZAUT, S. Cloning, characterization and expression of human and mouse lymphokine genes (interferon- γ and interleukin-2). International Lymphokine Workshop. Schloss Elmau (Germany), 1984.
- C.B.41 In Gory, C., Schiepl, A. and Landy, M. (eds.), "Cellular and Molecular Biology of Lymphokines", Academic Press, Inc., Orlando, pp. 385-401, 1985.
- C.B.42 FIEBS, W. The future role in medicine of proteins made by genetic engineering. Dehler Workshop on Biotechnology: Potentials and Limitations. Berlin (Germany), March 14-19, 1983.
- C.B.43 In Silver, G. (ed.), "Biotechnology: Potentials and Limitations", Dahlem Workshop Reports, Vol. 23, Springer Verlag, Berlin, pp. 165-177, 1984.
- C.B.44 FIEBS, W., BROUCKAERT, P., GUINET, Y., REZAUT, S., VAN ROY, P., DEVOS, R., FRANKEN, L., LECHEUX-ROELS, C., MARCENOT, A., TAVENIER, J., and VAN DER NIEUW, J. Recombinant Interferon γ and its synergism with tumor necrosis factor in the human and mouse systems. The 1985 WHO-15th Meeting on the Interferon System. Clearwater Beach (FL, USA), October 11-18, 1985.
- C.B.45 In Schellekens, M. and Stewart, L. W. (eds.), "The Biology of the Interferon System 1985", Elsevier Science Publishers, Amsterdam-New York-Oxford, pp. 241-248, 1986.
- C.B.46 FIEBS, W., BROUCKAERT, P., DEVOS, R., FRANKEN, L., LECHEUX-ROELS, C., REZAUT, S., SUFFIA, P., TAVENIER, J., VAN DER NIEUW, J., and VAN ROY, P. Lymphokines and cytokines in anti-cancer therapy. 51st Symposium on Quantitative Biology. Cold Spring Harbor (New York, USA), May 18 - June 4, 1986.
- C.B.47 In "Molecular Biology of Neoplasia", Cold Spring Harbor Symposium on Quantitative Biology, Vol. 51, Cold Spring Harbor Laboratory, Cold Spring Harbor, pp. 347-353, 1986.
- C.B.48 FIEBS, W. Van nucleotides in tumor. Voorrecht N.W. Beijerincklabor, Kon. Ned. Academie 21 Juni 1986.
- C.B.49 F., FIEBS, W., and COX, J. Induction of 35 kDa protein gp94 in human cells treated with recombinant human tumor necrosis factor (TNF- α). The 1986 WHO-15th Meeting on the Interferon System. Clearwater Beach (FL, USA), October 11-18, 1986.
- C.B.50 In Cantell, M. and Schellekens, M. (eds.), "The Biology of the Interferon System 1985", Martinus Nijhoff Publishers, Dordrecht-Sutton-Lancaster, pp. 211-221, 1985.
- C.B.51 REVILLACQUA, M. P., MILLER, M. K., PORTER, J. S., FIEBS, W., MENDRICK, D. L., CUTRAN, D. J., and CHENOMUTH, M. A. Endothelial-dependent mechanisms of leukocyte adhesion: Regulation by interleukin-1 and tumor necrosis factor. 4th International Congress on Immunology, Toronto (Canada), 1984.
- C.B.52 In Novel, H. E. (ed.), "Leukocyte Emigration and its Sequelae", Karger, Basel, pp. 38-51, 1981.
- C.B.53 FIEBS, W., BROUCKAERT, P., GOLDBERG, A., REITZEL, I., SUFFIA, P., TAVENIER, J., VANHAESBROECK, B., and VAN ROY, P. Structure-function relationship of tumor necrosis factor and its mechanism of action. Ciba Foundation Symposium No. 111: Tumor Necrosis Factor and Related Cytokines. London (United Kingdom), January 19-23, 1982.
- C.B.54 In Rock, C. and March, J. (eds.), "Tumor Necrosis Factor and Related Cytokines", John Wiley & Sons, Chichester, pp. 159-169, 1983.
- C.B.55 FIEBS, W., REZAUT, S., BROUCKAERT, P., EVERAERT, R., SUFFIA, P., TAVENIER, J., VANHAESBROECK, B., and VAN ROY, P. Tumor Necrosis Factor: A potential antitumor agent? J. Interferon Res. 7, 317-322, 1987.
- C.B.56 FIEBS, W., REZAUT, S., BROUCKAERT, P., EVERAERT, R., VANHAESBROECK, B., SUFFIA, P., TAVENIER, J., VANHAESBROECK, B., and VAN ROY, P. Gene cloning and structure-function relationship of cytokines such as TNF and Interleukin-1. Immunol. Lett. 16, 219-224, 1987.
- C.B.57 FIEBS, W., BROUCKAERT, P., DEVOS, R., FRANKEN, L., LECHEUX-ROELS, C., MARCENOT, A., REZAUT, S., SUFFIA, P., TAVENIER, J., and VAN ROY, P. Molecular and cellular biology of tumor necrosis factor: Interferon- γ and their synergism. The 1986 WHO-15th Meeting on the Interferon System (Clearwater Beach, FL, USA), September 7-13, 1986.
- C.B.58 In Cantell, M. and Schellekens, M. (eds.), "The Biology of the Interferon System 1985", Martinus Nijhoff Publishers, Dordrecht-Sutton-Lancaster, pp. 203-216, 1985.

- C.b.32 **PIER, V., DE WALLE, P., VAN DE VOORDE, A., CARMEL, P. and FIER, W.**
Expression of functional mouse antibodies directed against the tumour marker human placental alkaline phosphatase in non-lymphoid cells.
Int. J. Cancer 41 (Suppl. 2), 28-29, 1988.
- C.b.33 **PIER, W., BEIER, M., BROUCKERT, P., EVERAERT, B., KATZMAN, G., RUFFIN, P., TAVENIER, J., VANHAESBROECK, B. and VAN ROY, F.**
The potential use of an antitumour agent.
30th 1988 Congress on Cytokinines Laboratory and Clinical Evaluation. London (United Kingdom), October 19-21, 1987.
In Gearing, A.J.M. and Henderson, W. (eds.), "Development in Biological Standardization", Vol. 89, Sarger, Basel, pp. 111-121, 1988.
- C.b.34 **PIER, W., LEHMANS, R., MEYERS, M., KARLON, E., STAMMERS, P., STEINLE, L., VAN MECHELEN, E. and STAMUT, E.**
Engineering animal expression of heterologous genes in gram-negative bacteria.
8th International Biotechnology Symposium. Paris (France), July 17-21, 1988.
In Durand, G., Sobichon, L. and Fibrant, J. (eds.), "Proceedings", Vol. 3, Societe Francaise de Microbiologie, Paris, pp. 480-481, 1988.
- C.b.35 **MULLERBACH, D., MEYERS, G., FIER, W., STAMMERS, M., VULLER, A., LILJESTROM, P. and CAMER, M.**
Viral delivery systems for heterologous antigens and vaccines.
Virus Symposium, Park City (UT, USA), January 30 - February 6, 1988.
In Leach, L. (ed.), "Technological Advances in Vaccine Development", Alan R. Liss, Inc., New York, pp. 278-291, 1988.
- C.b.36 **KARIEL, M., COOPMAN, P., DRAGONETTI, C., FIER, W., GAO, J., MESSIAH, L. and VAN POI, P.**
Spontaneous and oncogene-mediated acquisition of the invasive phenotype by cells in culture.
In Ricolson, G.L. and Fidler, I.J. (eds.), "Tumor Progression and Metastasis", UCLA Symposium on Molecular and Cellular Biology, New Series, Vol. 18, Alan R. Liss, Inc., New York, pp. 119-128, 1988.
- C.b.37 **VAN DE VOORDE, A., PIER, V., DE WALLE, P., CARMEL, P. and FIER, W.**
Yellering of an anti-human placental alkaline phosphatase immunoglobulin using genetic engineering.
WATO Advanced Workshop, Brussels (Belgium), 1987.
In Miller, A.D.A. (ed.), "Advanced Research on Animal Cell Technology", Elsevier Academic Publishers, Amsterdam, pp. 233-248, 1988.
- C.b.38 **PIER, W., BROUCKERT, P., COMTE, J., COMTE, R., EVERAERT, B., GUERIN, Y., LILJESTROM, P., SPANGLER, D., TAKAHASHI, M., TIGNON, S., VANHAESBROECK, B. and VAN ROY, F.**
Interleukin-6: Biological function and regulation of the gene expression in vitro and in vivo.
6th International Congress on Immunopharmacology, Osaka (Japan), May 19-23, 1988.
In Hadden, J.W., Sprent, J., Yokoyama, Y., Austen, K.F., Dayer, P. and Mosch, M. (eds.), "Advances in Immunopharmacology", Vol. 2, Pergamon Press, Oxford, pp. 103-108, 1988.
- C.b.39 **PIER, W., BEIER, M., BROUCKERT, P., EVERAERT, B., CHOTIM, J., KATZMAN, G., LILJESTROM, P., TAKAHASHI, M., TAVENIER, J., VAN ROY, F., VANHAESBROECK, B., VAN VLIET, K. and VAN ROY, F.**
Tumour necrosis factor and interleukin-6: Structure and mechanism of action of the molecular, cellular and in vivo levels.
Symposium on Virus-cell interaction and vectors for transfer and propagation of genes, Maastricht (Netherlands), October 18-20 and 21-23, 1988.
In Leach, L., Dayer, P. and Osterley, M. (eds.), "Vectors as Tools for the Study of Normal and Abnormal Growth and Differentiation", NATO ASI Series, Vol. 18, Springer-Verlag, Berlin, pp. 219-224, 1988.
- C.b.40 **PIER, W.**
Biological properties of tumour necrosis factor.
International Conference on Growth Inhibitors: Pre-clinical and Clinical Evaluation in Cancer, Oxford (United Kingdom), March 20-22, 1988.
In "Documentation", IAC Technical Services, London, 1988.
- C.b.41 **PIER, W.**
Tumour necrosis factor: From the clone to the clinic.
Bouilles de la Science et des Technologies 7, 41-55, 1988.
- C.b.42 **PIER, W.**
The Mechanism of action in vitro and in vivo.
Poche Closed Symposium, Tokyo (Japan), April 10-12, 1988.
In Ueda, K., Futuhal, Y. and Breda, J. (eds.), "Influence of Molecular Biology on Drug Discovery", Clinical Pharmacology, Vol. 11, W. Bockmeyer Verlag, Munchen, pp. 17-18, 1988.

- Board Assigned Page #153

- C.B.17 BROUHAERT, P., FORNBERG, R., LIRINT, C.,
TAKAHASHI, W., CAMMEL, A. and FIERS, W.
Strategies to broaden the therapeutic margin of
Tumor Necrosis Factor
4th International Conference on Tumor Necrosis
Factor, Veldhoven (The Netherlands), May 2-8, 1991.
In Fiers, W. & Boudan, W.A. (eds.), "Tumor Necrosis
Factor: Molecular and Cellular Biology and Clinical
Relevance", Harger, Basel, pp. 218-222, 1991.
- C.B.18 VAN OUDK, X., TAYLOR, J. and FIERS, W.
Structure-activity studies of human Tumor Necrosis
Factor
Protein Engineering 1, 3-22, 1994.
- C.B.19 BIVATY, R. and FIERS, W.
Molecular mechanisms of tumor necrosis factor-
induced cytotoxicity: what we do understand and what
we do not.
TIBS Lett. 110, 9-14, 1994.
- C.B.20 BROUHAERT, P., AMELIOT, P., CAMMEL, A., FURNBERG,
R., LIRINT, C., TAKAHASHI, W., VAN HOLLE, B. and
FIERS, W.
Receptor-mediated mutants of tumor necrosis factor
in the therapy of cancer: Preclinical studies.
Circ. Shock 41, 183-189, 1991.
- C.B.21 VANDERVELLE, P., COOSENS, V., BIVATY, R.,
DECLERCQ, S., CROOTEN, J., VANHAESEBROECK, B., VAN
DE CRAEM, R., VERCAUTEN, D., DEPUTOT, B., DEMEYER,
G. and FIERS, W.
Functional requirement of the two TNF receptors for
induction of apoptosis in PCa cells and the role of
mitochondria in TNF-induced cytotoxicity.
Circ. Shock 46, 188-199, 1994.
- C.B.22 HARTMAN, G. and FIERS, W.
TNF-induced mechanisms for its gene induction.
In Fackel, L. & Mitt, R. (eds.), "Signalling
Mechanisms - From Transcription Factors to Oxidative
Stress", NATO ASI Series, Vol. H 32, Springer,
Berlin/Heidelberg/New York, pp. 133-142, 1993.
- C.B.23 VANDEWAELE, P., DECLERCQ, W., BIVATY, R. and
FIERS, W.
Two tumor necrosis factor receptors: structure and
function.
Trends Cell Biol. 3, 181-189, 1993.

C. GENERAL (MOLECULAR BIOLOGY, VIROLOGY, ETC.)

- C.G.1 FIERSE, W.
De chemische structuur van plantenvirussen.
Med. VI. Chem. Ver. 18, 138-139, 1956.
- C.G.2 FIERSE, W.
The primary structure of nucleic acids.
Revue Ferment. Industr. Affin. 18, 36-41, 1944.
- C.G.3 FIERSE, W.
De informatiecode der levende cel.
Med. VI. Chem. Ver. 18, 91-127, 1954.
- C.G.4 FIERSE, W.
Le cellule vivente doud le laboratoire chimique.
Chimie du 30e Siecle 17, 87-94, 1944.
- C.G.5 FIERSE, W.
R.C. Shorans. Modelprijze voor Geneeskunde 1948.
Med. VI. Chem. Ver. 16, 108-193, 1948.
- C.G.6 FIERSE, W.
Perspectieven in de moleculaire biologie.
Chem. Weekblad 88, 17-18, 1971.
- C.G.7 FIERSE, W.
Virussen: Algemeene Inleiding.
Verhand. Fac. Landbouwet. Rij 10, 1-22, 1971.
- C.G.8 FIERSE, W., SCHILL, J. and VAN MONTGOU, M.
De nieuwe mogelijkheden voor wetenschappelijk onderzoek.
Chemie Magazine 1, 13-15, 1974.
- C.G.9 FIERSE, W.
Gene.
In "McGraw-Hill Yearbook Science and Technology",
McGraw-Hill Book Co., New York, pp. 189-211, 1974.
- C.G.10 FIERSE, W.
Het gebruik van restrictie-enzymen bij de sequentie-analyse van DNA.
Chem. Weekblad 71, 9-12, 1973.
- C.G.11 VAN DE VOORDE, A. and FIERSE, W.
Bankerel-gerichte chemotherapie.
In "Reis en herber overvinnen", Belgisch Werk tegen
Banker, Brussel, pp. 27-29, 1981.
- C.G.12 CORTEBAST, R. and FIERSE, W.
Gene.
In "McGraw-Hill Encyclopedia of Science and
Technology", McGraw-Hill Book Company, New York, pp.
118-126, 1981.
- C.G.13 FIERSE, W.
Nieuwe mogelijkheden van de biotechnologie.
Corectie 4, 333-337, 1981.

D. ABSTRACTS

Total number: 302.


Dec 30 20 02

[Signature]

Commissioner of Patents
Commissaire des brevets

[Signature]

Examinateur



This is EXHIBIT FIERS-3
to
the Affidavit of Walter C. Fiers
sworn before me
this 13th day of November, 2001

Commissioner for Oath or Notary Public

Derynck lab

21/41

- Guanidiniumchloride . . . 4 ml gelatin of 1 ml Gd } RNA of 1

- Zinner IF $2-3 \times 10^8$ U / mg.

mens, mens -

mens: all fibrillat-type

Zinnering - polyD-agarose

affigel met zeifels ontvuld. → gel { 35 K
22 K

verlating 85% 35 K.

verit met actif biol. molecule.

langzet: 2% wavy - 2 differe banden
fingerprint getijl.

Zinner: portinisch ok
{ onticellulair actif.

- Spec. kolon: 1. Zee - delect of 1a - { Fil.
2. Blue Sefence. { 1a.

- Bolken. Kolon intinell: geen vertis v. sturist

- Kankor: 1. Kankor / met voor dufin.

- Glycolytic PAS (pratic uit - Sleff).

Reager 35 K +
22 K -

langzet: { neuromitose belanden → 1 band (IEF).

Cor A - column: nusselt 2 jalen -

Tunicamycin: niets mer verbanden → glycolytic } 1a
by glycolytic met vertis van lid act.

Anger zelfs heelt mer glycolytic nusselt volledig
te eliminer door tunicamycin belandling.

Intinell: vertis v. lid act v. Fil-IF
mer intin te beest.

- Inductie : poly I:C + HEAT-doxon \rightarrow gaat om nagen.


- Productie : Membran : veel ook fibr. 3-20%
Fibr. : " " leuk. - 1-20% (lang af v. menen v. infectie)

human interferon: specifiek.
gevoel v. stimulatie v. Thymus.

poly I:C + poly I:C : weinig nevensmerdingen - kan als behandeling aan patiënten toegepast worden.

- mRNA : translatie - mens \rightarrow X. leenis.

Revel : ook in retic. - minder 1/100



Cloning : 4 firma's Score 1
6 groepen -


- chromosom: receptor 21
productie: 2, 3 - ook van 21.

- receptors: heel actief. - binding van gebeld IF
moleculaire veranderingen in IF bevestigde cellen.

- anticellulair effect: neutrocyt celgroei - niet veranderig. morfologie
effect - G1.

- effect of tumoren: vermindering extracellulair virusel antige
vermindering intracellulair - defect of reactie

- Revel: effect of uncoupling?
IF 20u na infectie.




cellulose and vegetable.

$$111 A' 15' A 2' 5' A = 2' 5' A$$

IF + Lo RNA + ATP


2) lineare \rightarrow $\left\{ \begin{array}{l} \text{Erfaktor 2.} \\ \text{lineare} \end{array} \right.$



{ tRNA plebs & not onygar
 methyloster

2'5' A cell in normal culture - endogly 5x low IF
- clinical: L 1210 → L 1210 B (IF resistant - large receptor).
↓ ↓
tumor tumor IF
IF generally IF → cell culture - therapy resistant

mens



mens $2-5 \times 10^5$ U/kg.

meer trials, men 99% leucocytes -

+ Karyo bij oeg.

eventueel geen positief effect indien ook andere
behandeling (bv. steroïden!).

- Ebola

- Hepatitis B -

- Osteosarcoma

Standaard.

verlozing

15-65-75%

{ Steroïden kunnen mogelijk zijn

{ Niet of kleine hoeveelheden.

{ Voornamelijk meer sensitief.

Mamografie:

- IF - is equivalent aan leucocytes.

Fibrinolyse:

{ metastasiserende

{ hepatitis B

{ oeg hepatitis

{ fibrinolyse : geen level.

{ leucocytes : wel "

Neuromusculaire:

Leeds : onguisabelen?

via postganglionaire?

belang is bloedend (minder met PIF)

BIUGEN S.A.

Opened <u>Dec 30</u> 20 <u>02</u>
Déclaré(e) le <u>[Signature]</u>
Commissioner of Patents Commissaire des brevets
In presence of examiner <u>[Signature]</u> en présence de l'examineur

Reply To
Suite 3700
One New York Plaza
New York, New York 100.

December 1, 1978

Prof. Dr. W. Fiers
Laboratorium voor
Moleculaire Biologie
9000 Ghent
BELGIUM

Dear Dr. Fiers:


This will confirm our meeting at 8 o' clock a.m. in Ghent on Thursday, December 14. I will have with me Deborah Masters who is one of Biogen's vice presidents. We will meet you at the Ghent railway station. We need to take a 11 o' clock train from Ghent in order to catch the 11:43 a.m. train to Paris.

I look forward to seeing you on the 14th.

Sincerely,

[Signature]
Daniel D. Adams
President &
Managing Director

DDA:bjb



This is EXHIBIT FIER-4
to
the Affidavit of Walter C. Fiers
sworn before me
this 19th day of November, 2001

Commissioner for Oath or Notary Public

11b Avenue de la Porte-Neuve, Luxembourg

9/2/79

11²⁰

BIOGEN

SCIENTIFIC BOARD MTG. (PARIS)

Robin Nicholson

Ken Murray

Bernard Mach

Ray Schaeffer

Moshe Afi

Bob Luciano

Wally Gilbert

Charles Weissmann

Peter Hans Hofschneider

Heinz Schaller



Walter Fiers

Dag Laurason

Brian Hartley

Philippe Kourilsky

Phil Sharpe

Opened	Dec 30	20 02
Decachetée le		
		
Commissioner of Patent Commissaire aux brevets		
		
In presence of examiner en présence de l'examineur		



This is EXHIBIT FIER-5
to
the Affidavit of Walter C. Fiers
sworn before me
this 19th day of November, 2001

Commissioner for Oath or Notary Public

WALTER FIER

done in collaboration w/ 2 other
groups

1- in Lorraine

Desomer no longer active

exp. in assay for interf.
prepare cells - running bioassays

2- Comte

Posters Institute of Brussels
all translation work.

- Mol. Biol - my Fier's group in Ghent

Start of fibroblast RNA - gradient \rightarrow
partial purif. - $\sim 40 \times$ purif.

~ 150 clones most screening on this disc.
then 20,000 clones

Using dAT tails (consider
compared to C.W. work) - higher
level of trans formation - \therefore used this for
higher yields.

* Techniques / ^{cloning} same as Chas. Weissman

Detection & screening:
Hybrid Assisted Translation ---
wanted to be less dependent on
posit. artifacts.

System has not worked - despite
apparent functioning of controls -

Assays -

① Orig - same as C.W. -
same tech. by content in several

②: Reticuloendothelial
assay. biol. activity

Much more active in #1 than #2 (20x)

Screened 150 clones in groups of 50 -
mixed results.

HAT - -

out of protein = same whether
or not you melt it. - see that system works.

Satellite Tobacco Necrosis Virus
(STNV) is used as marker. (?)



Good anti-interferon / ~~also~~ also
Grip down ~~the~~ STNV.

very good messenger
in terms of activity

either way works or

interf. as induces in mysterious way

Screening 20,000 in groups of 50

Opened	Dec 30	2002
Détachée le		
		
Commissaire of Patents Commissaire des brevets		
In presence of 		
en présence de l'examineur		



This is EXHIBIT FIERS-6
 to
 the Affidavit of Walter C. Fiers
 sworn before me
 this 19th day of November, 2001

 Commissioner for Oath or Notary Public

- De positieve resultaten op DBM-culturen zijn zeer duidelijk, ook op herhaling (behalve voor B).

Gel ④ bevat ook kolom 13, die wegen twijfelachtig positief resultaat nu ook duidelijk ④.

- De keuze voor andere groyselectie valt op C.

- = ⊕ op cellulose (ex).
- = ⊕ op biologische activiteit. (plaque reduction assay is maar in verduidelijking tot gevolgt)
- = ⊕ op Sijthausse → andere afwijking
→ dit kolonijgebonden DNA is verder gewenst
gewenst over suikerefficientie
(op cellulose zit: bevat nog erg veel RNA)
- = geen enkele maar ⊖ gekooid!

C = 2'1/2 (A→D) - 46 kolonies

1. opgeven van tray 2'1/2 op dish - 10 dagen/bet.

2. enten van pre-kulturen LB/bet

$A_2 \rightarrow C_1$	$\left[\begin{array}{l} A_2 \rightarrow C_1 \\ C_2 \rightarrow D_{12} \end{array} \right]$
$C_2 \rightarrow D_{12}$	

3. enten van 400 ml kulturen LB/bet

$A_2 \rightarrow C_1$	$\left[\begin{array}{l} A_2 \rightarrow C_1 \\ C_2 \rightarrow D_{12} \end{array} \right]$	\rightarrow (tot 9.2.1)
$C_2 \rightarrow D_{12}$		

4. lyse-procedure

na afcentrifugeren vd kulturen (GS-3, 10 min - 3000 g) (3x per pot)
2x wasen met 1x TES


heropnemen in 100 ml suikrose 10% - Tris 50 mM pH 8
+ 10 ml lysozyme (50 mg/ml Tris 25 mM pH 8) (± 30 min)
+ 10 ml EDTA (10.5M pH 8)
+ 80 ml TLN (over de middag)

alles bij
keluitemperatuur.

Clearing spin : 6 poly-allonies (10 ml) gr groep (\rightarrow 3 groepen)
24 K - 45 min.

5. PEG.-precipitatie : SN van Clearing spin + 1/3 vol 40% PEG
2M NaCl

overnacht in ijskoude kamer (ert of yn)



Het inviezen nl pellets (na opgroeien) is wel de oorzaak van de aanwezigheid van RNA in de DNA-preparaten na lye en Cell gradient-centrifugatie !

C - subgroepen

2' - 12

C₁: A₂ → A₉ (8)

C₂: A₁₀ → A₁₁, B₁ → B₆ (8)

C₃: B₇ → B₁₂, C₁ (7)

C₄: C₂ → C₉ (8)

C₅: C₁₀ → C₁₂, D₁ → D₅ (8)

C₆: D₆ → D₁₂ (7)

7. CCl₄-vermischings-gradiëntcentrifugatie (2 per groep, totaal 12)

RGOT:
40 K
13°C
over weekend.

→ banden goed te herkennen - onderdeel van Rv17

8. Aftappen met equivalente punke → 5 ml tevens spijt

IAA extractie:
+ 3 vol H₂O
+ 1/10 vol NaAc 2M pH 5.1
+ 2 vol EtOH

overnacht -20°C

9. EtOH petten afdrogen


heropnemen in 1/10 STE
fenolwateren - fenolwater warmen
etherextractie 3x
ether verdampen

tot vol per groep = 1 ml 1/10 STE

↓
2 µl op gel laden (1% agarose)
+ 1 µg pBR 322 referentie

10. Sukrose-methylegradiënt-centrifugatie

↓
wegen afwezig zijn de RNA
⇒ NIET NODIG



SUBGROEPEN C

1 2 3 4 5 6

↓
pBR-322 referentie

⇒ (0.3) (0.3) (0.4) ± 1 µg (1) (0.1) (0.2)

$$O_2 : A_{10} \rightarrow A_{11}, B_1 \rightarrow B_6 \quad (6)$$



$$O_3 : B_1 \rightarrow B_{12}, C_1 \quad (7)$$

$$O_4 : C_2 \rightarrow C_9 \quad (8)$$

$$O_5 : C_{12} \rightarrow C_{12}, D_1 \rightarrow D_5 \quad (8)$$

$$O_6 : D_6 \rightarrow D_{12} \quad (7)$$



Rezultati na kolonijupridrobu



SUBGROEPEN ○

1 2 3 4 5 6


U₁ (+)



→ 14pg 13 19 (10) 12 13 10 13 (10pg)

Er is een weinig RNA aanwezig (→ 3+4 scores geven een idee, 5 en 6 afleest echter niet meer)

3,4,5 bevatten chromosomaal DNA.



0₁ 0₂ 0₃ 0₄ 0₅ 0₆

gekineerd STNV-RNA

→ 7M NaOH-gel

[3,4. → jantel-banden , STNV-RNA na CIP-behandeling.
(calf-internu-phosphatex)
5,6,7,8,9 → jantel-banden , STNV-RNA , zonder voorbehandeling.
(In gescelte-volgorde).

22/11

RNA bereidingen (Rege - Leiden)

- $\left\{ \begin{array}{lll} W_1: & \text{non induced} & \text{VGS \# 23, 10 cellen} & (10/10/79) \\ W_2: & \text{induced} & \text{VGS \# 24, 20 cellen} & (24/10/79) \\ W_3: & \text{induced} & \text{VGS \# 26, 15 cellen} & (16/11/79). \end{array} \right.$

→ overgebruikt als EtOH precipitaat.

* afcentrifugeren 9000 tpm HB4 -20°C

heropnemen TE + 0.1% SDS. 400 µl

+ 1 vol CHCl_3 / IAA

+ 1 vol fenol

en fenolfase naar boven met 50 µl TE

SN 400 + 50 µl pook en naar boven met 1 vol CHCl_3 / IAA
(verwijderen van fenol)

* EtOH precipiteren

↓

22/11

heropnemen in 80 µl TE, 0.1% SDS.

↓ -20°C

2 µl staalje, OD meten.


W_1	0.101	$\times 2 \times 100 \rightarrow$	20.2 OD/ml	$\times 0.6 \rightarrow$	12.12 OD	$\times 40$	485
W_2	0.144		28.8 OD/ml		17.28 OD		691
W_3	0.150		30.0 OD/ml		18.00 OD		720

indemerkend in batches

- $\left\{ \begin{array}{l} W_1: 3 \times 170 \mu\text{g} \\ W_2: 4 \times 20 \mu\text{g} + 4 \times 115 \mu\text{g} \\ W_3: 4 \times 30 \mu\text{g} + 4 \times 132 \mu\text{g} \end{array} \right.$

→ alles EtOH precipiteren i.o.v. 0.27 l. tot pH 5.1

1 Methode 1



10/4/00

DSM - cellulose ~ DNA

hybridisatie met plopp
0.05%

hybridisatie overnacht


daarna 7x wassen

3x elutie

alles present behandelen in effluentie / centrifugatie

Dec 30 2002
Opened / Declassified
STNY-RVA
Commissioner of Patents
Commissaire des brevets
In presence of / en présence de l'examinateur

Dus per hybridisatie hebben we




hierbij is FT: droog
W: was
el: eluatie

Alle fracties worden nu na samen voegen en vóór ethanol precipitatie samengevoegd vandelid in 1 hel of ten

(A) 1/2 → precipitatie met halfloos cRNA

(B) 1/2 → " poly A⁺ RNA / terugbrengen



This is EXHIBIT FIERS-7
to
the Affidavit of Walter C. Fiers
sworn before me
this 19th day of November, 2001

W. L. S. M. S.

voor beiden onderzoek kunnen passen C en O in aanpak, op

[illegible]

19/10/85

Gelatin was in all the specimens from the program
from 1864. About a liter in P.P.
of an old program specimen taken in at
London, program was well kept
but it did not contain any and was
the opposite of the old.

From our search the program and
the kind of it in P.P. was not at
all kept (I know not, it is
a program) and the program is 1864 line
man and kind of it in the
specimens taken.

Besluit

C₃ zijn product van 26K en reactief van biolakt I.

100

Q. 1. 2. 3. 4.

O_2 is negativ von 26K zu positiv von 26K zu 27K

Co ook psem another banden (#264) to see
ma zetic + I know. p.ac. and ant. IF

O₂ sub pump

Biologische Aktivität

clone	op DBP-cell. pouder		op DBP-cell. fitter		op mlu-collabo fitter	
	FT	SL	FT	SL	FT	SL
O ₂ 12	< 0.2	0	0.7	0	2.0	0
2	3.2	0	0.2	0	0.7	0
3	2.2	0	2.8	1.2	2.0	0
4	2.2	0	1.0	0	2.2	0
5	0.7	0	0.7	50.2	2.2	0
6	0.7	0	2.0	50.2	0.5	0
7	0.5	0	2.2	0	2.0	0.5
8			50.2	2.2	2.0	0.7
9						
O ₂ unmutated	2.2	0.5	50.2	2.2		

Dem O₂ 12 is duidelijk probeel voor IF-aktiviteit.

Dec 30 02
[Signature]
M. G. J. J. J.
en présence de l'examinateur



This is EXHIBIT FIERS-8
to
the Affidavit of Walter C. Fiers
sworn before me
this 19th day of November, 2001

Commissioner for Oath or Notary Public

after SDS-polyacrylamide gel electrophoresis, the eluate should be centrifuged at 20,000 rev/min (Sorvall SS-34 rotor) for 20 minutes to remove particulate matter before dialysis. Coomassie blue staining of the gels to locate protein bands does not interfere with subsequent sequenator analysis.

New technologies such as the improved amino acid sequencing method described above lead to new research opportunities. With the greater sensitivity provided by this technique, we now can obtain amino acid sequence information on both proteins and peptides with submicrogram (picomole) quantities. This sensitivity should permit analysis of biomedically relevant molecules—such as the interferons—that can only be obtained in microgram quantities, and this ability opens possibilities for further study of these molecules. For example, knowledge of the amino acid sequence permits the synthesis of corresponding DNA probes and opens the possibility of

new strategies for isolating genes, such as those for interferons, that express low levels of messenger RNA's (8).

MICHAEL W. HUNKAPILLER

LEROY E. HOOD

Division of Biology, California Institute of Technology, Pasadena 91125

References and Notes

1. A. Isaacs and J. Lindenmann, *Proc. R. Soc. London Ser. B* 147, 358 (1957).
2. T. Menges, *N. Engl. J. Med.* 300, 42 (1979).
3. E. Knight, Jr., M. W. Hunkapiller, B. D. Korant, R. W. F. Hardy, L. E. Hood, *Science* 207, 525 (1980); K. C. Zoon, M. E. Smith, P. J. Bridges, C. B. Anshen, M. W. Hunkapiller, L. E. Hood, *ibid.*, p. 527; M. Taira, R. J. Brown, B. M. Javaram, P. Lengyel, M. W. Hunkapiller, L. E. Hood, *ibid.*, p. 528.
4. P. Edman and G. Berg, *Eur. J. Biochem.* 1, 80 (1967).
5. For a review of these methods, see M. Hunkapiller and L. Hood, *Biochemistry* 17, 2124 (1978).
6. B. Wittmann-Liebold, *Hoppe-Seyler's Z. Physiol. Chem.* 354, 1413 (1973).
7. N. Johnson, M. Hunkapiller, L. Hood, *Anal. Biochem.* 100, 335 (1979).
8. B. Noyes, M. Taira, R. Stein, E. Agarwal, *Proc. Natl. Acad. Sci. U.S.A.* 76, 1770 (1979).
9. Supported by a gift from the Ben Weingart Foundation.

29 November 1979

Human Fibroblast Interferon: Amino Acid Analysis and Amino Terminal Amino Acid Sequence

Abstract. The purification of human fibroblast interferon has been simplified to a two-step procedure consisting of affinity chromatography on Blue Sepharose and sodium dodecyl sulfate polyacrylamide gel electrophoresis. A preliminary amino acid composition and the sequence of the 13 amino-terminal residues of homogeneous interferon prepared by this method is reported.

Since the discovery of interferon, its purification and chemical characterization have been primary goals of interferon research. Although their attainment has been slow because of the small quantities of interferon proteins avail-

able, purification to homogeneity has now been achieved with some interferons. However, only microgram quantities have been available for characterization—human fibroblast interferon (1, 2), human lymphoblastoid interferon (3), human leukocyte interferon (4), mouse interferon (5)—and only limited structural information has been acquired (4, 6).

A thorough understanding at the molecular level of the numerous phenomena that are caused by interferon in cells in culture and in animals will not be possible until the elucidation of primary and secondary structures of the interferon proteins is achieved. This structural information will permit (i) comparison of amino acid sequences of interferons from various cell types and animal species, (ii) identification of the polypeptide segments involved in binding to interferon-specific cell-surface receptors, and (iii) chemical synthesis of interferons.

We now report an improved procedure for the purification of human fibroblast interferon that can be used to provide enough protein for structural studies.

Using the automated protein microsequencing technique described in (7), we have determined the sequence of the 13 amino acid residues at the amino terminus of the interferon prepared by this method. We also report a preliminary amino acid composition of the protein.

Human diploid fibroblast cells (FS-4) were cultured and interferon was produced (1). Interferon was assayed by a microtechnique (8) with vesicular stomatitis virus as the challenge virus. Interferon units are given in National Institutes of Health human fibroblast interferon units.

The crude interferon, 10 to 15 liters produced in the absence of serum, was made 1M in NaCl and passed at room temperature through a column (4 by 10 cm) of Blue Sepharose (Pharmacia, Inc.) equilibrated with 0.02M sodium phosphate buffer, pH 7.2, containing 1M NaCl. The interferon was retained whereas more than 95 percent of the total protein passed through the column. The interferon was eluted with a mixture of the column buffer and ethylene glycol (1:1), and each fraction was diluted immediately with 0.5 volume of the buffer (Fig. 1a). Fractions containing interferon activity were pooled, diluted with two volumes of the column buffer, and passed through a small (1 by 6 cm) Blue Sepharose column for concentration. The interferon was eluted as described above (Fig. 1b).

Fractions containing interferon were pooled, dialyzed against 1 mM Tris-HCl,




Fig. 1. (a) Fractionation of crude interferon on a large column of Blue Sepharose. Elution of interferon with 50 percent ethylene glycol in column buffer begins at fraction 1. (b) Small Blue Sepharose column. Fractions 7 to 17 in (a) were pooled, passed through the small column, and eluted with 50 percent ethylene glycol in column buffer (fractions 1 to 20).

Table 1. Amino acid composition of human fibroblast interferon.

Amino acid	Composition	
	Mole percent	Residues per 20,000 daltons
Asp	11.3	18.9
Thr	4.8	6.8
Ser	6.2	10.5
Glu	15.9	27.0
Pro	1.6	2.7
Gly*	4.6	7.8
Ala	5.9	10.0
Cys†	1.0	1.7
Val	3.3	6.0
Met	1.7	2.9
Ile	5.3	9.0
Leu	12.0	20.4
Tyr	4.4	7.5
Phe	5.5	9.4
His	2.9	4.9
Lys	6.8	11.6
Arg	6.4	10.9
Trp‡	0.6	1.0

*Includes correction for free glycine present in unhydrolyzed protein. †Determined after performic acid oxidation. ‡Determined after hydrolysis with mercaptoethanesulfonic acid.




Fig. 2. (a) Preparative electrophoresis of interferon activity profile. Fractions 8 to 13 in Fig. 1b were pooled, concentrated, and subjected to electrophoresis in a polyacrylamide slab gel, 0.75 mm thick. Fractions 14 and 15 were pooled and processed for amino acid sequencing. (b) Polyacrylamide slab gel, staining of proteins eluted from preparative gel in (a). Approximately 2 percent of the protein in fractions 14 and 15 (a) was subjected to electrophoresis and stained. Lanes 1 and 3, standard proteins; lane 2, interferon.

pH 6.8, containing 0.02 percent sodium dodecyl sulfate (SDS, Bio-Rad electrophoresis grade), and concentrated to dryness in a vacuum centrifuge. The interferon was then subjected to electrophoresis on a SDS-polyacrylamide slab gel and eluted (Fig. 2a). Fractions eluted from the gel were assayed for interferon activity (Fig. 2a). Approximately 0.2 μ g of interferon from the peak activity fraction was subjected to electrophoresis in this system again, and the gel was stained with Coomassie blue (Fig. 2b).

The preparative electrophoresis fractions containing interferon were pooled and centrifuged for 30 minutes at 30,000 rev/min at 4°C to remove polyacrylamide gel particles. The interferon solution was dialyzed first against 0.15M NaCl containing 0.1 percent SDS and then against 0.02 percent SDS. The dialyzed interferon was concentrated to dryness in a vacuum centrifuge.

This purification procedure is simpler and shorter than that described previously (1). Recoveries from the large Blue Sepharose column have ranged from 50 to 100 percent, and those from the small column approach 100 percent. The interferon (5×10^5 U/mg) eluted from these columns is stable for at least 4 weeks at 4°C in 1M NaCl, 35 percent ethylene glycol, pH 7.2. Recoveries of activity from the SDS gels have ranged from 5 to 20 percent, and specific activities of this protein have ranged from 2×10^5 to 8×10^5 U/mg. Accurate specific activities are difficult to determine, and




Fig. 3. The amino-terminal amino acid sequence of human fibroblast interferon.

two- to fourfold differences above 1×10^5 U/mg are probably not meaningful. Overall yields of purified interferon from 10- to 15-liter batches of crude material (5×10^5 to 7×10^5 total units, 8×10^5 U/mg) have averaged around 10 percent. This gives 5 to 10 μ g of homogeneous interferon.

Amino acid analysis on 1- to 2- μ g portions was performed on a Durrum D-500 amino acid analyzer (Table 1). Automated Edman degradation on 0.4- to 2- μ g portions of the purified interferon was performed on a spinning cup sequencer (7). Phenylthiohydantoin (PTH) amino acids were identified by high-performance liquid chromatography (HPLC) on a Du Pont Zorbax CN column (9).

The sequence of the 13 amino terminal amino acid residues of human fibroblast interferon was determined by this microsequencing technique (Fig. 3). Yields of PTH methionine at cycle 1 for three sequencer runs ranged from 60 to 100 percent (based on protein determination by amino acid analysis), and the sequencer repetitive cycle yields were 92 to 95 percent. Any unblocked minor peptide sequence present at > 5 percent of the reported sequence could have been detected by the methods used, but none has

homogeneity of the interferon peptide preparation.

Determining the amino acid sequence of a protein is essential in order to identify its active site and to understand its molecular mechanism of action. Comparison of structural features of interferons from different species and from different cell types within an animal will prove or disprove whether they are different proteins. If there is an active site common to all interferons, it should be identifiable by comparison of the amino acid sequences. Comparison of the amino terminal sequence reported here for human fibroblast interferon does not as yet reveal any apparent homology with the amino-terminal sequence reported for human lymphoblastoid interferon (16). However, there is limited homology (3/13 residues identical) with the 37,000 dalton mouse Ehrlich ascites cell interferon (11).

E. KNIGHT, J.

Central Research & Development
Department, E. I. du Pont
de Nemours & Company,
Wilmington, Delaware 19898

M. W. HUNKAPILLER

Division of Biology, California Institute
of Technology, Pasadena 91109

B. D. KORANT, R. W. F. HODGSON
Central Research & Development
Department, E. I. du Pont
de Nemours & Company

L. E. HOOD

Division of Biology,
California Institute of Technology

References and Notes

1. E. Knight, Jr., *Proc. Natl. Acad. Sci. U.S.A.* **73**, 520 (1976).
2. W. Berthold, C. Tan, Y. H. Tan, *J. Biol. Chem.* **253**, 5206 (1978).
3. K. C. Zoon, M. E. Smith, P. J. Bridgen, D. J. Nedden, C. B. Anfinsen, *Proc. Natl. Acad. Sci. U.S.A.*, in press.
4. M. Rubinstein, S. Rubinstein, P. C. Fanillet, R. S. Miller, A. A. Walman, S. Pestka, *ibid.* **7**, 643 (1979).
5. E. Knight, Jr., *J. Biol. Chem.* **250**, 4139 (1975); Y. Yamamoto and Y. Kawade, *J. Gen. Virol.* **33**, 225 (1976); M. Kawada, B. Cabrer, J. Taura, M. Rebellato, E. Statter, H. Weidlich, *Langmuir, J. Biol. Chem.* **254**, 598 (1979); J. C. Mueser-Guignard, M. G. Tavery, J. Gresser, J. De Maeyer, *Nature (London)* **271**, 622 (1978).
6. B. Cabrer, H. Taura, R. J. Broeze, T. D. Kemp, K. Williams, E. Statter, W. H. Kunitzberg, *Langmuir, J. Biol. Chem.* **254**, 3681 (1979); Y. F. Tan, F. Haraoka, W. Berthold, H. Smith-Hanzen, C. Tan, *ibid.*, p. 8067.
7. M. W. Hunkapiller and L. E. Hood, *Science* **207**, 523 (1980).
8. J. A. Armstrong, *Appl. Microbiol.* **21**, 71 (1971).
9. N. J. Johnson, M. W. Hunkapiller, L. E. Hood, *Anal. Biochem.*, in press.
10. K. C. Zoon, M. E. Smith, P. J. Bridgen, C. B. Anfinsen, M. W. Hunkapiller, L. E. Hood, *Science* **207**, 527 (1980).
11. H. Taura, R. J. Broeze, B. M. Jayaram, J. Langmuir, M. W. Hunkapiller, L. E. Hood, *ibid.* p. 328.

30 November 1979

SCIENCE, VOL. 207